

BAG4 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16332b**Specification**

BAG4 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O95429
Other Accession	Q8CI61 , NP_004865.1
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	49594
Antigen Region	381-409

BAG4 Antibody (C-term) - Additional Information**Gene ID** 9530**Other Names**

BAG family molecular chaperone regulator 4, BAG-4, Bcl-2-associated athanogene 4, Silencer of death domains, BAG4, SODD

Target/Specificity

This BAG4 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 381-409 amino acids from the C-terminal region of human BAG4.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BAG4 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

BAG4 Antibody (C-term) - Protein Information**Name** BAG4

Synonyms SODD

Function Inhibits the chaperone activity of HSP70/HSC70 by promoting substrate release (By similarity). Prevents constitutive TNFRSF1A signaling. Negative regulator of PRKN translocation to damaged mitochondria.

Cellular Location

Cytoplasm.


Tissue Location

Ubiquitous.

BAG4 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

BAG4 Antibody (C-term) - Images

BAG4 Antibody (C-term) (Cat. #AP16332b) western blot analysis in HeLa cell line lysates (35ug/lane). This demonstrates the BAG4 antibody detected the BAG4 protein (arrow).

BAG4 Antibody (C-term) - Background

BAG4 is a member of the BAG1-related protein family. BAG1 is an anti-apoptotic protein that functions through interactions with a variety of cell apoptosis and growth related proteins including BCL-2, Raf-protein kinase, steroid hormone receptors, growth factor receptors and members of

the heat shock protein 70 kDa family. This protein contains a BAG domain near the C-terminus, which could bind and inhibit the chaperone activity of Hsc70/Hsp70. This protein was found to be associated with the death domain of tumor necrosis factor receptor type 1 (TNF-R1) and death receptor-3 (DR3), and thereby negatively regulates downstream cell death signaling. The regulatory role of this protein in cell death was demonstrated in epithelial cells which undergo apoptosis while integrin mediated matrix contacts are lost.

BAG4 Antibody (C-term) - References

Bailey, S.D., et al. Diabetes Care (2010) In press :
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Tao, H.F., et al. Zhongguo Shi Yan Xue Ye Xue Za Zhi 15(3):501-505(2007)
Riley, B.M., et al. Am. J. Med. Genet. A 143A (8), 846-852 (2007) :
Yang, Z.Q., et al. Cancer Res. 66(24):11632-11643(2006)