

# HTR7 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16398b

# **Specification**

# HTR7 Antibody (C-term) - Product Information

Application WB,E
Primary Accession P34969

Other Accession <u>NP\_062873.1</u>, <u>NP\_000863.1</u>

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
Catoliated MW
53555
Antigen Region

# HTR7 Antibody (C-term) - Additional Information

#### **Gene ID 3363**

### **Other Names**

5-hydroxytryptamine receptor 7, 5-HT-7, 5-HT-X, Serotonin receptor 7, HTR7

### Target/Specificity

This HTR7 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 405-433 amino acids from the C-terminal region of human HTR7.

### **Dilution**

WB~~1:1000

# **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

HTR7 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# HTR7 Antibody (C-term) - Protein Information

# Name HTR7

**Function** This is one of the several different receptors for 5- hydroxytryptamine (serotonin), a biogenic hormone that functions as a neurotransmitter, a hormone, and a mitogen. The activity of



this receptor is mediated by G proteins that stimulate adenylate cyclase.

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein.

### **Tissue Location**


Isoform A is the predominant isoform in spleen, caudate and hippocampus. Isoform B is expressed at lower levels Isoform D is a minor isoform in terms of expression

## HTR7 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# HTR7 Antibody (C-term) - Images



HTR7 Antibody (C-term) (Cat. #AP16398b) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the HTR7 antibody detected the HTR7 protein (arrow).

# HTR7 Antibody (C-term) - Background

The neurotransmitter, serotonin, is thought to play a role in various cognitive and behavioral functions. The serotonin receptor encoded by this gene belongs to the superfamily of G protein-coupled receptors and the gene is a candidate locus for involvement in autistic disorder and other neuropsychiatric disorders. Three splice variants have been identified which encode proteins that differ in the length of their carboxy terminal ends.

# HTR7 Antibody (C-term) - References

Ruano, G., et al. Pharmacogenomics 11(7):959-971(2010)





Tel: 858.875.1900 Fax: 858.875.1999

Pinheiro, A.P., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (5), 1070-1080 (2010): Corominas, R., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (1), 177-184 (2010): Wei, Z., et al. Prog. Neuropsychopharmacol. Biol. Psychiatry 33(3):547-551(2009) Iceta, R., et al. J. Physiol. Pharmacol. 60(1):157-164(2009)