

NAT5 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16414b

Specification

NAT5 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P61599
Other Accession	Q7ZXR3 , P61600 , Q2PFM2 , Q58ED9 , NP_852669.1 , NP_852668.1 , NP_057184.1
Reactivity	Human
Predicted	Zebrafish, Monkey, Mouse, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	20368
Antigen Region	127-155

NAT5 Antibody (C-term) - Additional Information

Gene ID 51126

Other Names

N-alpha-acetyltransferase 20, Methionine N-acetyltransferase, N-acetyltransferase 5, N-terminal acetyltransferase B complex catalytic subunit NAA20, N-terminal acetyltransferase B complex catalytic subunit NAT5, NatB complex subunit NAT5, NatB catalytic subunit, NAA20, NAT5

Target/Specificity

This NAT5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 127-155 amino acids from the C-terminal region of human NAT5.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NAT5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

NAT5 Antibody (C-term) - Protein Information

Name NAA20

Synonyms NAT5

Function Catalytic subunit of the NatB complex which catalyzes acetylation of the N-terminal methionine residues of peptides beginning with Met-Asp, Met-Glu, Met-Asn and Met-Gln (PubMed:34230638). Proteins with cell cycle functions are overrepresented in the pool of NatB substrates. Required for maintaining the structure and function of actomyosin fibers and for proper cellular migration.

Cellular Location


Cytoplasm. Nucleus

NAT5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NAT5 Antibody (C-term) - Images



NAT5 Antibody (C-term) (Cat. #AP16414b) western blot analysis in A2058 cell line lysates (35ug/lane). This demonstrates the NAT5 antibody detected the NAT5 protein (arrow).

NAT5 Antibody (C-term) - Background

NAT5 is a component of N-acetyltransferase complex B (NatB). Human NatB performs cotranslational N(alpha)-terminal acetylation of methionine residues when they are followed by asparagine (Starheim et al., 2008 [PubMed 18570629]). [supplied by OMIM].

NAT5 Antibody (C-term) - References

Kuo, H.P., et al. Sci Signal 3 (108), RA9 (2010) :
Evjenth, R., et al. J. Biol. Chem. 284(45):31122-31129(2009)
Polevoda, B., et al. BMC Proc 3 SUPPL 6, S2 (2009) :
Starheim, K.K., et al. Biochem. J. 415(2):325-331(2008)
Sanchez-Puig, N., et al. Protein Sci. 15(8):1968-1976(2006)