

RBM26 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16434a**Specification**

RBM26 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q5T8P6
Other Accession	Q6NZN0 , NP_071401.3
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	113597
Antigen Region	196-225

RBM26 Antibody (N-term) - Additional Information**Gene ID** 64062**Other Names**

RNA-binding protein 26, CTCL tumor antigen se70-2, RNA-binding motif protein 26, RBM26, C13orf10

Target/Specificity

This RBM26 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 196-225 amino acids from the N-terminal region of human RBM26.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RBM26 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RBM26 Antibody (N-term) - Protein Information**Name** RBM26 ([HGNC:20327](#))

Synonyms C13orf10


Function May be involved in the turnover of nuclear polyadenylated (pA+) RNA.

RBM26 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RBM26 Antibody (N-term) - Images



RBM26 Antibody (N-term) (Cat. #AP16434a) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the RBM26 antibody detected the RBM26 protein (arrow).

RBM26 Antibody (N-term) - Background

RBM26 (RNA binding motif protein 26) bears two RNA recognition motif (RRM) domains and a C3H1-type zinc finger. The function remains unknown.

RBM26 Antibody (N-term) - References

Rose, J. Phd, et al. Mol. Med. (2010) In press :
Miele, A., et al. J. Cell. Biochem. 102(1):136-148(2007)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Olsen, J.V., et al. Cell 127(3):635-648(2006)
Beausoleil, S.A., et al. Nat. Biotechnol. 24(10):1285-1292(2006)