

**MANF Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP16437a****Specification**

---

**MANF Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P55145</a>
Other Accession	<a href="#">P0C5H9</a> , <a href="#">Q9CXI5</a> , <a href="#">P80513</a> , <a href="#">NP_006001.3</a>
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	20700
Antigen Region	23-51

**MANF Antibody (N-term) - Additional Information****Gene ID** 7873**Other Names**

Mesencephalic astrocyte-derived neurotrophic factor, Arginine-rich protein, Protein ARMET, MANF, ARMET, ARP

**Target/Specificity**

This MANF antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 23-51 amino acids from the N-terminal region of human MANF.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

MANF Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**MANF Antibody (N-term) - Protein Information****Name** MANF ([HGNC:15461](#))

**Synonyms** ARMET, ARP

**Function** Selectively promotes the survival of dopaminergic neurons of the ventral mid-brain (PubMed:[12794311](#)). Modulates GABAergic transmission to the dopaminergic neurons of the substantia nigra (By similarity). Enhances spontaneous, as well as evoked, GABAergic inhibitory postsynaptic currents in dopaminergic neurons (By similarity). Inhibits cell proliferation and endoplasmic reticulum (ER) stress-induced cell death (PubMed:[18561914](#), PubMed:[22637475](#), PubMed:[29497057](#)). Retained in the ER/sarcoplasmic reticulum (SR) through association with the endoplasmic reticulum chaperone protein HSPA5 under normal conditions (PubMed:[22637475](#)). Up-regulated and secreted by the ER/SR in response to ER stress and hypoxia (PubMed:[22637475](#)). Following secretion by the ER/SR, directly binds to 3-O-sulfogalactosylceramide, a lipid sulfatide in the outer cell membrane of target cells (PubMed:[29497057](#)). Sulfatide binding promotes its cellular uptake by endocytosis, and is required for its role in alleviating ER stress and cell toxicity under hypoxic and ER stress conditions (PubMed:[29497057](#)).


**Cellular Location**

Secreted. Endoplasmic reticulum lumen. Sarcoplasmic reticulum lumen. Note=Retained in the endoplasmic reticulum (ER), and sarcoplasmic reticulum (SR) under normal conditions (PubMed:[22637475](#)). Up-regulated and secreted by the ER/SR in response to ER stress and hypoxia (PubMed:[22637475](#), PubMed:[29497057](#))

**MANF Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**MANF Antibody (N-term) - Images**

MANF Antibody (N-term) (Cat. #AP16437a) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the MANF antibody detected the MANF protein (arrow).

**MANF Antibody (N-term) - Background**

Selectively promotes the survival of dopaminergic neurons of the ventral mid-brain. Modulates GABAergic transmission to the dopaminergic neurons of the substantia nigra. Enhances spontaneous, as well as evoked, GABAergic inhibitory postsynaptic currents in dopaminergic neurons (By similarity). Inhibits cell proliferation and endoplasmic reticulum (ER) stress-induced cell death.