

CEP72 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16441c

Specification

CEP72 Antibody (Center) - Product Information

Application WB,E **Primary Accession** O9P209 Other Accession NP 060610.2 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 71718 Antigen Region 231-259

CEP72 Antibody (Center) - Additional Information

Gene ID 55722

Other Names

Centrosomal protein of 72 kDa, Cep72, CEP72, KIAA1519

Target/Specificity

This CEP72 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 231-259 amino acids from the Central region of human CEP72.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CEP72 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

CEP72 Antibody (Center) - Protein Information

Name CEP72

Synonyms KIAA1519





Function Involved in the recruitment of key centrosomal proteins to the centrosome. Provides centrosomal microtubule-nucleation activity on the gamma-tubulin ring complexes (gamma-TuRCs) and has critical roles in forming a focused bipolar spindle, which is needed for proper tension generation between sister chromatids. Required for localization of KIZ, AKAP9 and gamma-tubulin ring complexes (gamma-TuRCs) (PubMed: 19536135). Involved in centriole duplication. Required for CDK5RAP22, CEP152, WDR62 and CEP63 centrosomal localization and promotes the centrosomal localization of CDK2 (PubMed: 26297806).

Cellular Location


Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite. Note=Localizes to the centrosome and centrosome-surrounding particles throughout the cell cycle. These particles disappear after microtubules are depolymerized using nocodazole, suggesting that CEP72-associating particles localize in a microtubule- dependent manner

CEP72 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CEP72 Antibody (Center) - Images




CEP72 Antibody (Center) (Cat. #AP16441c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the CEP72 antibody detected the CEP72 protein (arrow).

CEP72 Antibody (Center) - Background

CEP72 is a member of the

leucine-rich-repeat (LRR) superfamily of proteins. The protein is localized to the centrosome, a non-membraneous organelle that functions as the major microtubule-organizing center in animal cells.





CEP72 Antibody (Center) - References

McGovern, D.P., et al. Nat. Genet. 42(4):332-337(2010) Briggs, F.B., et al. Genes Immun. 11(3):199-208(2010) Oshimori, N., et al. EMBO J. 28(14):2066-2076(2009) Xin, X., et al. Genome Res. 19(7):1262-1269(2009) Olsen, J.V., et al. Cell 127(3):635-648(2006)