

**UHRF1BP1L Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP16540C****Specification**

---

**UHRF1BP1L Antibody (Center) - Product Information**

|                   |                             |
|-------------------|-----------------------------|
| Application       | WB,E                        |
| Primary Accession | <a href="#">A0JNW5</a>      |
| Other Accession   | <a href="#">NP_055869.1</a> |
| Reactivity        | Human                       |
| Host              | Rabbit                      |
| Clonality         | Polyclonal                  |
| Isotype           | Rabbit IgG                  |
| Calculated MW     | 164199                      |
| Antigen Region    | 544-572                     |

**UHRF1BP1L Antibody (Center) - Additional Information****Gene ID** 23074**Other Names**

UHRF1-binding protein 1-like, UHRF1BP1L, KIAA0701

**Target/Specificity**

This UHRF1BP1L antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 544-572 amino acids from the Central region of human UHRF1BP1L.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

UHRF1BP1L Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**UHRF1BP1L Antibody (Center) - Protein Information****Name** BLTP3B ([HGNC:29102](#))**Function** Tube-forming lipid transport protein which mediates the transfer of lipids between

membranes at organelle contact sites (PubMed:[35499567](#)). Required for retrograde traffic of vesicle clusters in the early endocytic pathway to the Golgi complex (PubMed:[20163565](#), PubMed:[35499567](#)).

#### Cellular Location


Cytoplasm, cytosol. Early endosome. Note=Localizes on a subpopulation of vesicle clusters in the early endocytic pathway

#### UHRF1BP1L Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### UHRF1BP1L Antibody (Center) - Images



UHRF1BP1L Antibody (Center) (Cat. #AP16540c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the UHRF1BP1L antibody detected the UHRF1BP1L protein (arrow).

#### UHRF1BP1L Antibody (Center) - References

Olsen, J.V., et al. Cell 127(3):635-648(2006)  
Strausberg, R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903(2002)  
Ishikawa, K., et al. DNA Res. 5(3):169-176(1998)