

ARNT2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16544c

Specification

ARNT2 Antibody (Center) - Product Information

Application IF, WB,E Primary Accession Q9HBZ2

Other Accession <u>Q78E60</u>, <u>Q61324</u>, <u>NP_055677.3</u>

Reactivity
Predicted
Mouse, Rat
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Mouse, Rat
Rabbit
Rabbit
Polyclonal
Rabbit IgG
78691
249-278

ARNT2 Antibody (Center) - Additional Information

Gene ID 9915

Other Names

Aryl hydrocarbon receptor nuclear translocator 2, ARNT protein 2, Class E basic helix-loop-helix protein 1, bHLHe1, ARNT2, BHLHE1, KIAA0307

Target/Specificity

This ARNT2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 249-278 amino acids from the Central region of human ARNT2.

Dilution

IF~~1:10~50 WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ARNT2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ARNT2 Antibody (Center) - Protein Information



Name ARNT2

Synonyms BHLHE1, KIAA0307

Function Transcription factor that plays a role in the development of the hypothalamo-pituitary axis, postnatal brain growth, and visual and renal function (PubMed:24022475). Specifically recognizes the xenobiotic response element (XRE).

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00981, ECO:0000269|PubMed:24465693}

ARNT2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

ARNT2 Antibody (Center) - Images



Fluorescent confocal image of SK-BR-3 cell stained with ARNT2 Antibody (Center)(Cat#AP16544c). SK-BR-3 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with ARNT2 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C). Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C). Nuclei were counterstained with DAPI (blue) (10 μ g/ml, 10 min).ARNT2 immunoreactivity is localized to nucleus significantly.





ARNT2 Antibody (Center) (Cat. #AP16544c) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the ARNT2 antibody detected the ARNT2 protein (arrow).

ARNT2 Antibody (Center) - Background

This gene encodes a member of the basic-helix-loop-helix-Per-Arnt-Sim (bHLH-PAS) superfamily of transcription factors. The encoded protein acts as a partner for several sensor proteins of the bHLH-PAS family, forming heterodimers with the sensor proteins that bind regulatory DNA sequences in genes responsive to developmental and environmental stimuli. Under hypoxic conditions, the encoded protein complexes with hypoxia-inducible factor 1alpha in the nucleus and this complex binds to hypoxia-responsive elements in enhancers and promoters of oxygen-responsive genes. A highly similar protein in mouse forms functional complexes with both aryl hydrocarbon receptors and Single-minded proteins, suggesting addition roles for the encoded protein in the metabolism of xenobiotic compounds and the regulation of neurogenesis, respectively.

ARNT2 Antibody (Center) - References

Ramirez, J.M., et al. Eur. J. Immunol. 40(9):2450-2459(2010) Jugessur, A., et al. PLoS ONE 5 (7), E11493 (2010) : Chakrabarti, B., et al. Autism Res 2(3):157-177(2009) Ferreira, M.A., et al. Nat. Genet. 40(9):1056-1058(2008) Martinez, V., et al. Breast Cancer Res. Treat. 110(3):521-530(2008)