

RYBP Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16629A**Specification**

RYBP Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q8N488
Other Accession	Q8CCI5 , NP_036366.3
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	24822
Antigen Region	1-30

RYBP Antibody (N-term) - Additional Information**Gene ID** 23429**Other Names**

RING1 and YY1-binding protein, Apoptin-associating protein 1, APAP-1, Death effector domain-associated factor, DED-associated factor, YY1 and E4TF1-associated factor 1, RYBP, DEDAF, YEAF1

Target/Specificity

This RYBP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human RYBP.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RYBP Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

RYBP Antibody (N-term) - Protein Information**Name** RYBP

Synonyms DEDAF, YEAF1

Function Component of a Polycomb group (PcG) multiprotein PRC1-like complex, a complex class required to maintain the transcriptionally repressive state of many genes, including Hox genes, throughout development. PcG PRC1-like complex acts via chromatin remodeling and modification of histones; it mediates monoubiquitination of histone H2A 'Lys-119', rendering chromatin heritably changed in its expressibility (PubMed:[25519132](#)). Component of a PRC1-like complex that mediates monoubiquitination of histone H2A 'Lys-119' on the X chromosome and is required for normal silencing of one copy of the X chromosome in XX females. May stimulate ubiquitination of histone H2A 'Lys-119' by recruiting the complex to target sites (By similarity). Inhibits ubiquitination and subsequent degradation of TP53, and thereby plays a role in regulating transcription of TP53 target genes (PubMed:[19098711](#)). May also regulate the ubiquitin-mediated proteasomal degradation of other proteins like FANK1 to regulate apoptosis (PubMed:[14765135](#), PubMed:[27060496](#)). May be implicated in the regulation of the transcription as a repressor of the transcriptional activity of E4TF1 (PubMed:[11953439](#)). May bind to DNA (By similarity). May play a role in the repression of tumor growth and metastasis in breast cancer by down-regulating SRRM3 (PubMed:[27748911](#)).

Cellular Location

Nucleus. Cytoplasm. Nucleus, nucleoplasm {ECO:0000250|UniProtKB:Q8CCI5}. Note=Primarily found in the nucleus Detected in a punctate pattern likely to represent Polycomb group (PcG) bodies (By similarity). {ECO:0000250|UniProtKB:Q8CCI5}

Tissue Location


Down-regulated in breast cancer tissues and in several breast cancer cell lines (at protein level) (PubMed:27748911) Widely expressed with highest levels in lymphoid tissues and placenta

RYBP Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RYBP Antibody (N-term) - Images



RYBP Antibody (N-term) (Cat. #AP16629a) western blot analysis in A375 cell line lysates (35ug/lane). This demonstrates the RYBP antibody detected the RYBP protein (arrow).



Western blot analysis of RYBP (arrow) using rabbit polyclonal RYBP Antibody (N-term) (Cat. #AP16629a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the RYBP gene.

RYBP Antibody (N-term) - Background

RYBP may be implicated in the regulation of the transcription as a repressor of the transcriptional activity of E4TF1. In tumor cell lines, may induce apoptosis.

RYBP Antibody (N-term) - References

- Rose, J. Phd, et al. Mol. Med. (2010) In press :
- Chen, D., et al. EMBO Rep. 10(2):166-172(2009)
- Novak, R.L., et al. Cancer Gene Ther. 15(11):713-722(2008)
- Danen-van Oorschot, A.A., et al. Cell Death Differ. 11(5):564-573(2004)
- Guelen, L., et al. Oncogene 23(5):1153-1165(2004)