

SEPT2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16761b

Specification

SEPT2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q15019
Other Accession	Q91Y81 , P42208 , Q2NKY7 , NP_001008492.1 , NP_001008491.1
Reactivity	Human
Predicted	Bovine, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	41487
Antigen Region	280-309

SEPT2 Antibody (C-term) - Additional Information

Gene ID 4735

Other Names

Septin-2, Neural precursor cell expressed developmentally down-regulated protein 5, NEDD-5, SEPT2, DIFF6, KIAA0158, NEDD5

Target/Specificity

This SEPT2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 280-309 amino acids from the C-terminal region of human SEPT2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SEPT2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SEPT2 Antibody (C-term) - Protein Information

Name SEPTIN2 ([HGNC:7729](#))

Function Filament-forming cytoskeletal GTPase. Forms a filamentous structure with SEPTIN12, SEPTIN6, SEPTIN2 and probably SEPTIN4 at the sperm annulus which is required for the structural integrity and motility of the sperm tail during postmeiotic differentiation (PubMed:[25588830](#)). Required for normal organization of the actin cytoskeleton. Plays a role in the biogenesis of polarized columnar-shaped epithelium by maintaining polyglutamylated microtubules, thus facilitating efficient vesicle transport, and by impeding MAP4 binding to tubulin. Required for the progression through mitosis. Forms a scaffold at the midplane of the mitotic spindle required to maintain CENPE localization at kinetochores and consequently chromosome congression. During anaphase, may be required for chromosome segregation and spindle elongation. Plays a role in ciliogenesis and collective cell movements. In cilia, required for the integrity of the diffusion barrier at the base of the primary cilium that prevents diffusion of transmembrane proteins between the cilia and plasma membranes: probably acts by regulating the assembly of the tectonic-like complex (also named B9 complex) by localizing TMEM231 protein. May play a role in the internalization of 2 intracellular microbial pathogens, *Listeria monocytogenes* and *Shigella flexneri*.

Cellular Location

Cytoplasm. Cytoplasm, cytoskeleton. Cytoplasm, cytoskeleton, spindle. Chromosome, centromere, kinetochore. Cleavage furrow. Midbody Cytoplasm, cell cortex. Cell projection, cilium membrane. Cell projection, cilium, flagellum. Note=In metaphase cells, localized within the microtubule spindle. At the metaphase plate, in close apposition to the kinetochores of the congressed chromosomes. In cells undergoing cytokinesis, localized to the midbody, the ingressing cleavage furrow, and the central spindle. During bacterial infection, displays a collar shape structure next to actin at the pole of invading bacteria. In epithelial cells, colocalizes with polyglutamylated tubulin around the trans-Golgi network, as well as juxatnuclear and proximal Golgi apparatus. Localizes at the base of the cilia near the morphological distinction between the cilia and plasma membranes. Found in the sperm annulus (PubMed:[25588830](#)).

Tissue Location

Widely expressed. Up-regulated in liver cancer.


SEPT2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SEPT2 Antibody (C-term) - Images





SEPT2 Antibody (C-term) (Cat. #AP16761b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the SEPT2 antibody detected the SEPT2 protein (arrow).



Western blot analysis of SEPT2 (arrow) using rabbit polyclonal SEPT2 Antibody (C-term) (Cat. #AP16761b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the SEPT2 gene.

SEPT2 Antibody (C-term) - Background

Filament-forming cytoskeletal GTPase. Required for normal organization of the actin cytoskeleton. Plays a role in the biogenesis of polarized columnar-shaped epithelium by maintaining polyglutamylated microtubules, thus facilitating efficient vesicle transport, and by impeding MAP4 binding to tubulin. Required for the progression through mitosis. Forms a scaffold at the midplane of the mitotic spindle required to maintain CENPE localization at kinetochores and consequently chromosome congression. During anaphase, may be required for chromosome segregation and spindle elongation. May play a role in the internalization of 2 intracellular microbial pathogens, *Listeria monocytogenes* and *Shigella flexneri*.

SEPT2 Antibody (C-term) - References

- Martins-de-Souza, D., et al. J Psychiatr Res (2010) In press :
- Guey, L.T., et al. Eur. Urol. 57(2):283-292(2010)
- Hosgood, H.D. III, et al. Respir Med 103(12):1866-1870(2009)
- Shen, M., et al. Environ. Mol. Mutagen. 50(4):285-290(2009)
- Mostowy, S., et al. J. Biol. Chem. 284(17):11613-11621(2009)