

**TTC8 Antibody (N-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP16824a**

**Specification**

**TTC8 Antibody (N-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">Q8TAM2</a>
Other Accession	<a href="#">NP_938051.1</a> , <a href="#">NP_653197.2</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	61534
Antigen Region	18-46

**TTC8 Antibody (N-term) - Additional Information**

**Gene ID** 123016

**Other Names**

Tetratricopeptide repeat protein 8, TPR repeat protein 8, Bardet-Biedl syndrome 8 protein, TTC8, BBS8

**Target/Specificity**

This TTC8 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 18-46 amino acids from the N-terminal region of human TTC8.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

TTC8 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**TTC8 Antibody (N-term) - Protein Information**

**Name** TTC8

## Synonyms BBS8

**Function** The BBSome complex is thought to function as a coat complex required for sorting of specific membrane proteins to the primary cilia. The BBSome complex is required for ciliogenesis but is dispensable for centriolar satellite function. This ciliogenic function is mediated in part by the Rab8 GDP/GTP exchange factor, which localizes to the basal body and contacts the BBSome. Rab8(GTP) enters the primary cilium and promotes extension of the ciliary membrane. Firstly the BBSome associates with the ciliary membrane and binds to RAB3IP/Rabin8, the guanosyl exchange factor (GEF) for Rab8 and then the Rab8-GTP localizes to the cilium and promotes docking and fusion of carrier vesicles to the base of the ciliary membrane. The BBSome complex, together with the LTZL1, controls SMO ciliary trafficking and contributes to the sonic hedgehog (SHH) pathway regulation. Required for proper BBSome complex assembly and its ciliary localization.

## Cellular Location

Cytoplasm, cytoskeleton, microtubule organizing center, centrosome. Cell projection, cilium membrane. Cytoplasm. Cytoplasm, cytoskeleton, microtubule organizing center, centrosome, centriolar satellite. Cell projection, cilium {ECO:0000250|UniProtKB:Q8VD72}

## Tissue Location


Widely expressed.

## TTC8 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## TTC8 Antibody (N-term) - Images



TTC8 Antibody (N-term) (Cat. #AP16824a) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the TTC8 antibody detected the TTC8 protein (arrow).

## TTC8 Antibody (N-term) - Background

This gene encodes a protein that has been directly linked to Bardet-Biedl syndrome. The primary features of this syndrome include retinal dystrophy, obesity, polydactyly, renal abnormalities and learning disabilities. Experimentation in non-human eukaryotes suggests that this gene is expressed in ciliated cells and that it is involved in the formation of cilia. Alternate transcriptional splice variants have been characterized.

#### **TTC8 Antibody (N-term) - References**

Riazuddin, S.A., et al. Am. J. Hum. Genet. 86(5):805-812(2010)  
Bin, J., et al. Hum. Mutat. 30 (7), E737-E746 (2009) :  
Chung, W.K., et al. Hum. Hered. 67(3):193-205(2009)  
Nachury, M.V., et al. Cell 129(6):1201-1213(2007)  
Ansley, S.J., et al. Nature 425(6958):628-633(2003)