

## TRPC1 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP16863b

## **Specification**

## TRPC1 Antibody (C-term) - Product Information

Application WB,E
Primary Accession P48995

Other Accession <u>Q9TUN9</u>, <u>Q61056</u>, <u>Q18784</u>, <u>NP 003295.1</u>

Reactivity Human

Predicted Bovine, Mouse, Rabbit

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 91212
Antigen Region 765-793

# TRPC1 Antibody (C-term) - Additional Information

#### **Gene ID 7220**

### **Other Names**

Short transient receptor potential channel 1, TrpC1, Transient receptor protein 1, TRP-1, TRPC1, TRP1

## Target/Specificity

This TRPC1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 765-793 amino acids from the C-terminal region of human TRPC1.

#### **Dilution**

WB~~1:1000

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

TRPC1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

# TRPC1 Antibody (C-term) - Protein Information

#### Name TRPC1





## Synonyms TRP1

**Function** Forms a receptor-activated non-selective calcium permeant cation channel. Probably is operated by a phosphatidylinositol second messenger system activated by receptor tyrosine kinases or G-protein coupled receptors. Also activated by intracellular calcium store depletion.

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein

## **Tissue Location**


Seems to be ubiquitous.

## TRPC1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# TRPC1 Antibody (C-term) - Images



TRPC1 Antibody (C-term) (Cat. #AP16863b) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the TRPC1 antibody detected the TRPC1 protein (arrow).





Anti-TRPC1 Antibody (C-term) at 1:1000 dilution + HepG2 whole cell lysate Lysates/proteins at 20  $\mu$ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 91 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

# TRPC1 Antibody (C-term) - Background

TRPC1 belongs to the transient receptor potential (TRP) superfamily of cation channels. TRP cation channels are involved in diverse physiologic processes, including receptor- and store-operated Ca(2+) entry, mineral absorption, and cell death. They also function as sensors for pain, heat, cold, sound, stretch, and osmotic changes (Zhang et al., 2009 [PubMed 19193631]).

# TRPC1 Antibody (C-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010) Bomben, V.C., et al. Glia 58(10):1145-1156(2010) Lu, M., et al. J. Mol. Endocrinol. 44(5):285-294(2010) Gonzalez-Cobos, J.C., et al. Front. Biosci. 15, 1023-1039 (2010): Ingueneau, C., et al. J. Cell. Mol. Med. 13 (8B), 1620-1631 (2009):