

CSPG5 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16908b**Specification**

CSPG5 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O95196
Other Accession	NP_006565.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	60016
Antigen Region	530-559

CSPG5 Antibody (C-term) - Additional Information**Gene ID** 10675**Other Names**

Chondroitin sulfate proteoglycan 5, Acidic leucine-rich EGF-like domain-containing brain protein, Neuroglycan C, CSPG5, CALEB, NGC

Target/Specificity

This CSPG5 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 530-559 amino acids from the C-terminal region of human CSPG5.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CSPG5 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CSPG5 Antibody (C-term) - Protein Information**Name** CSPG5

Synonyms CALEB, NGC

Function May function as a growth and differentiation factor involved in neuritogenesis. May induce ERBB3 activation.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q9ERQ6}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q9ERQ6} Synaptic cell membrane {ECO:0000250|UniProtKB:Q71M36}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q71M36}. Endoplasmic reticulum membrane {ECO:0000250|UniProtKB:Q71M36}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q71M36}. Golgi apparatus membrane {ECO:0000250|UniProtKB:Q71M36}; Single-pass type I membrane protein {ECO:0000250|UniProtKB:Q71M36}. Cell surface {ECO:0000250|UniProtKB:Q71M36}. Secreted. Note=In neurons, localizes to synaptic junctions. Also detected in the endoplasmic reticulum and the Golgi Partially enriched in lipid rafts. {ECO:0000250|UniProtKB:Q71M36, ECO:0000250|UniProtKB:Q9ERQ6}


Tissue Location

Detected in cerebrospinal fluid (at protein level) (PubMed:25326458). Detected in urine (at protein level) (PubMed:37453717). Expressed in brain (at protein level) (PubMed:9950058).

CSPG5 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CSPG5 Antibody (C-term) - Images

CSPG5 Antibody (C-term) (Cat. #AP16908b) western blot analysis in U251 cell line lysates

(35ug/lane). This demonstrates the CSPG5 antibody detected the CSPG5 protein (arrow).

CSPG5 Antibody (C-term) - Background

CSPG5 may function as a growth and differentiation factor involved in neuritogenesis. May induce ERBB3 activation.

CSPG5 Antibody (C-term) - References

So, H.C., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 153B (1), 103-113 (2010) :
Ichihara-Tanaka, K., et al. J. Biol. Chem. 281(41):30857-30864(2006)
Aono, S., et al. J. Neurosci. Res. 83(1):110-118(2006)
Kinugasa, Y., et al. Biochem. Biophys. Res. Commun. 321(4):1045-1049(2004)
Hassel, B., et al. J. Biol. Chem. 278(41):40136-40143(2003)