

F9 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP16976c

Specification

F9 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	P00740
Other Accession	NP_000124.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	51778
Antigen Region	266-295

F9 Antibody (Center) - Additional Information

Gene ID 2158

Other Names

Coagulation factor IX, Christmas factor, Plasma thromboplastin component, PTC, Coagulation factor IXa light chain, Coagulation factor IXa heavy chain, F9

Target/Specificity

This F9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 266-295 amino acids from the Central region of human F9.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

F9 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

F9 Antibody (Center) - Protein Information

Name F9

Function Factor IX is a vitamin K-dependent plasma protein that participates in the intrinsic pathway of blood coagulation by converting factor X to its active form in the presence of Ca(2+) ions, phospholipids, and factor VIIIa (PubMed:[8295821](#), PubMed:[2592373](#), PubMed:[20121197](#), PubMed:[20121198](#), PubMed:[1730085](#), PubMed:[19846852](#), PubMed:[39880037](#)).

Cellular Location

Secreted

Tissue Location


Detected in blood plasma (at protein level) (PubMed:19846852, PubMed:2592373, PubMed:3857619, PubMed:8295821, PubMed:9169594). Synthesized primarily in the liver and secreted in plasma.

F9 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

F9 Antibody (Center) - Images



F9 Antibody (Center) (Cat. #AP16976c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the F9 antibody detected the F9 protein (arrow).



All lanes : Anti-F9 Antibody (Center) at 1:2000 dilution Lane 1: 293T/17 whole cell lysate Lane 2: A431 whole cell lysate Lane 3: HepG2 whole cell lysate Lane 4: K562 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 51 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

F9 Antibody (Center) - Background

This gene encodes vitamin K-dependent coagulation factor IX that circulates in the blood as an inactive zymogen. This factor is converted to an active form by factor X α , which excises the activation peptide and thus generates a heavy chain and a light chain held together by one or more disulfide bonds. The role of this activated factor IX in the blood coagulation cascade is to activate factor X to its active form through interactions with Ca $^{+2}$ ions, membrane phospholipids, and factor VIII. Alterations of this gene, including point mutations, insertions and deletions, cause factor IX deficiency, which is a recessive X-linked disorder, also called hemophilia B or Christmas disease.

F9 Antibody (Center) - References

Bailey, S.D., et al. *Diabetes Care* 33(10):2250-2253(2010)
Yang, L., et al. *J. Biol. Chem.* 285(37):28488-28495(2010)
Kao, C.Y., et al. *Thromb. Haemost.* 104(2):355-365(2010)
Roberts, K.E., et al. *Gastroenterology* 139(1):130-139(2010)
Arellano, A.R., et al. *J. Thromb. Haemost.* 8(5):1132-1134(2010)