

VDAC3 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17092c

Specification

VDAC3 Antibody (Center) - Product Information

Application	WB, IHC-P-Leica, IHC-P, FC,E
Primary Accession	Q9Y277
Other Accession	Q9TT13 , Q29380 , Q60931 , Q9MZ13 , NP_005653.3 , NP_001129166.1
Reactivity	Human, Mouse, Rat
Predicted	Bovine, Pig, Rabbit
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	156-183

VDAC3 Antibody (Center) - Additional Information

Gene ID 7419

Other Names

Voltage-dependent anion-selective channel protein 3, VDAC-3, hVDAC3, Outer mitochondrial membrane protein porin 3, VDAC3

Target/Specificity

This VDAC3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 156-183 amino acids from the Central region of human VDAC3.

Dilution

WB~~1:1000
IHC-P-Leica~~1:500
IHC-P~~1:100
FC~~1:25

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

VDAC3 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

VDAC3 Antibody (Center) - Protein Information

Name VDAC3 ([HGNC:12674](#))

Function Non-selective voltage-gated ion channel that mediates the transport of anions and cations through the mitochondrion outer membrane and plasma membrane (PubMed:[31935282](#)). Forms a high-conducting channel with a stable open state and a voltage-induced closure with a mild preference for anions over cations (PubMed:[31935282](#)). Involved in male fertility and sperm mitochondrial sheath formation (By similarity).

Cellular Location

Mitochondrion outer membrane {ECO:0000250|UniProtKB:P21796}. Membrane Note=May localize to non-mitochondrial membranes

Tissue Location


Expressed in erythrocytes (at protein level) (PubMed:27641616). Widely expressed. Highest in testis (PubMed:9781040).

VDAC3 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


VDAC3 Antibody (Center) - Images




VDAC3 Antibody (Center) (Cat. #AP17092c) western blot analysis in NCI-H460 cell line lysates (35ug/lane). This demonstrates the VDAC3 antibody detected the VDAC3 protein (arrow).




All lanes : Anti-VDAC3 Antibody (Center) at 1:1000 dilution Lane 1: Human heart lysate Lane 2: NCI-H460 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: U-2OS whole cell lysate Lane 5: Mouse testis lysate Lane 6: Rat brain lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




All lanes : Anti-VDAC3 Antibody (Center) at 1:1000 dilution Lane 1: Human heart lysate Lane 2: NCI-H460 whole cell lysate Lane 3: A431 whole cell lysate Lane 4: U-2OS whole cell lysate Lane 5: Mouse testis lysate Lane 6: Mouse heart lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 31 kDa Blocking/Dilution buffer: 5% NFDM/TBST.




Immunohistochemical analysis of paraffin-embedded human testis tissue using AP17092c performed on the Leica® BOND RXm. Samples were incubated with primary antibody(1/500) for 1 hours at room temperature. A undiluted biotinylated CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of AP17092c on paraffin-embedded Human heart tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of AP17092c on paraffin-embedded Human testis tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Overlay histogram showing Hela cells stained with AP17092c(green line). The cells were fixed with 2% paraformaldehyde (10 min) and then permeabilized with 90% methanol for 10 min. The cells were then incubated in 2% bovine serum albumin to block non-specific protein-protein interactions followed by the antibody (AP17092c, 1:25 dilution) for 60 min at 37°C. The secondary antibody used was Goat-Anti-Rabbit IgG, DyLight® 488 Conjugated Highly Cross-Adsorbed(1583138) at 1/200 dilution for 40 min at 37°C. Isotype control antibody (blue line) was rabbit IgG1 (1µg/1x10⁶ cells) used under the same conditions. Acquisition of >10, 000 events was performed.

VDAC3 Antibody (Center) - Background

VDAC3 belongs to a group of mitochondrial membrane channels involved in translocation of adenine nucleotides through the outer membrane. These channels may also function as a mitochondrial binding site for hexokinase (see HK1; MIM 142600) and glycerol kinase (GK; MIM 300474) (Rahmani et al., 1998).[supplied by OMIM].

VDAC3 Antibody (Center) - References

Reina, S., et al. FEBS Lett. 584(13):2837-2844(2010)

Lefievre, L., et al. Proteomics 7(17):3066-3084(2007)

Lamesch, P., et al. Genomics 89(3):307-315(2007)

Ewing, R.M., et al. Mol. Syst. Biol. 3, 89 (2007) :

Rush, J., et al. Nat. Biotechnol. 23(1):94-101(2005)

VDAC3 Antibody (Center) - Citations

- [Changes in the mitochondrial protein profile due to ROS eruption during ageing of elm \(*Ulmus pumila* L.\) seeds.](#)