

CHRNG Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17105a

Specification

CHRNG Antibody (N-term) - Product Information

WB,E **Application Primary Accession** P07510 NP 005190.4 Other Accession Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 57883 Antigen Region 22-50

CHRNG Antibody (N-term) - Additional Information

Gene ID 1146

Other Names

Acetylcholine receptor subunit gamma, CHRNG, ACHRG

Target/Specificity

This CHRNG antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 22-50 amino acids from the N-terminal region of human CHRNG.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CHRNG Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CHRNG Antibody (N-term) - Protein Information

Name CHRNG (HGNC:1967)

Synonyms ACHRG





Tel: 858.875.1900 Fax: 858.875.1999

Function After binding acetylcholine, the AChR responds by an extensive change in conformation that affects all subunits and leads to opening of an ion-conducting channel across the plasma membrane.

Cellular Location


Postsynaptic cell membrane; Multi-pass membrane protein. Cell membrane; Multi-pass membrane protein

CHRNG Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

CHRNG Antibody (N-term) - Images



CHRNG Antibody (N-term) (Cat. #AP17105a) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the CHRNG antibody detected the CHRNG protein (arrow).

CHRNG Antibody (N-term) - Background

The mammalian muscle-type acetylcholine receptor is a transmembrane pentameric glycoprotein with two alpha subunits, one beta, one delta, and one epsilon (in adult skeletal muscle) or gamma (in fetal and denervated muscle) subunit. This gene, which encodes the gamma subunit, is expressed prior to the thirty-third week of gestation in humans. The gamma subunit of the acetylcholine receptor plays a role in neuromuscular organogenesis and ligand binding and disruption of gamma subunit expression prevents the correct localization of the receptor in cell membranes. Mutations in this gene cause Escobar syndrome and a lethal form of multiple pterygium syndrome. Muscle-type acetylcholine receptor is the major antigen in the autoimmune disease myasthenia gravis.





CHRNG Antibody (N-term) - References

Saccone, N.L., et al. Genes Brain Behav. (2010) In press:
Gratacos, M., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 150B (6), 808-816 (2009):
Saccone, N.L., et al. Am. J. Med. Genet. B Neuropsychiatr. Genet. 150B (4), 453-466 (2009):
Zouridakis, M., et al. Biochim. Biophys. Acta 1794(2):355-366(2009)
Chang, B., et al. Int. J. Surg. Pathol. 17(1):6-15(2009)