

## CDS2 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17134c

## Specification

# CDS2 Antibody (Center) - Product Information

Application Primary Accession Other Accession Reactivity Predicted Host Clonality Isotype Calculated MW Antigen Region WB,E <u>O95674</u> <u>O91XU8, O99L43, A0JNC1, NP\_003809.1</u> Human Bovine, Mouse, Rat Rabbit Polyclonal Rabbit IgG 51418 138-165

# CDS2 Antibody (Center) - Additional Information

Gene ID 8760

**Other Names** 

Phosphatidate cytidylyltransferase 2, CDP-DAG synthase 2, CDP-DG synthase 2, CDP-diacylglycerol synthase 2, CDS 2, CDP-diglyceride pyrophosphorylase 2, CDP-diglyceride synthase 2, CTP:phosphatidate cytidylyltransferase 2, CDS2

## Target/Specificity

This CDS2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 138-165 amino acids from the Central region of human CDS2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### Precautions

CDS2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## CDS2 Antibody (Center) - Protein Information



# Name CDS2 (HGNC:1801)

**Function** Catalyzes the conversion of phosphatidic acid (PA) to CDP- diacylglycerol (CDP-DAG), an essential intermediate in the synthesis of phosphatidylglycerol, cardiolipin and phosphatidylinositol (PubMed:<u>25375833</u>). Exhibits specificity for the nature of the acyl chains at the sn-1 and sn-2 positions in the substrate, PA and the preferred acyl chain composition is 1-stearoyl-2-arachidonoyl-sn- phosphatidic acid (PubMed:<u>25375833</u>). Plays an important role in regulating the growth and maturation of lipid droplets which are storage organelles at the center of lipid and energy homeostasis (PubMed:<u>26946540</u>, PubMed:<u>31548309</u>).

#### **Cellular Location**

Endoplasmic reticulum membrane; Multi-pass membrane protein

#### **Tissue Location**

Widely expressed. Expressed in heart, brain and retina, and to a lesser extent in placenta, lung, liver, skeletal muscle, kidney and pancreas.

# **CDS2 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# CDS2 Antibody (Center) - Images



CDS2 Antibody (Center) (Cat. #AP17134c) western blot analysis in HepG2 cell line lysates (35ug/lane).This demonstrates the CDS2 antibody detected the CDS2 protein (arrow).

## CDS2 Antibody (Center) - Background



Breakdown products of phosphoinositides are ubiquitous second messengers that function downstream of many G protein-coupled receptors and tyrosine kinases regulating cell growth, calcium metabolism, and protein kinase C activity. This gene encodes an enzyme which regulates the amount of phosphatidylinositol available for signaling by catalyzing the conversion of phosphatidic acid to CDP-diacylglycerol. This enzyme is an integral membrane protein localized to two subcellular domains, the matrix side of the inner mitochondrial membrane where it is thought to be involved in the synthesis of phosphatidylglycerol and cardiolipin and the cytoplasmic side of the endoplasmic reticulum where it functions in phosphatidylinositol biosynthesis. Two genes encoding this enzyme have been identified in humans, one mapping to human chromosome 4q21 and a second to 20p13.

# CDS2 Antibody (Center) - References

Bailey, S.D., et al. Diabetes Care (2010) In press : Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) : Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009) Olsen, J.V., et al. Cell 127(3):635-648(2006) Olsen, J.V., et al. Cell 127(3):635-648(2006)