

**ATP2C1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP17155b****Specification**

---

**ATP2C1 Antibody (C-term) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P98194</a>
Other Accession	<a href="#">NP_001001486.1</a> , <a href="#">NP_001001485.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	100577
Antigen Region	882-909

**ATP2C1 Antibody (C-term) - Additional Information****Gene ID** 27032**Other Names**

Calcium-transporting ATPase type 2C member 1, ATPase 2C1, ATP-dependent Ca(2+) pump PMR1, ATP2C1, KIAA1347, PMR1L

**Target/Specificity**

This ATP2C1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 882-909 amino acids from the C-terminal region of human ATP2C1.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

ATP2C1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**ATP2C1 Antibody (C-term) - Protein Information****Name** ATP2C1 {ECO:0000303|PubMed:10615129, ECO:0000312|HGNC:HGNC:13211}**Function** ATP-driven pump that supplies the Golgi apparatus with Ca(2+) and Mn(2+) ions, both

essential cofactors for processing and trafficking of newly synthesized proteins in the secretory pathway (PubMed:[16192278](#), PubMed:[30923126](#), PubMed:[21187401](#), PubMed:[12707275](#), PubMed:[20439740](#)). Within a catalytic cycle, acquires  $\text{Ca}(2+)$  or  $\text{Mn}(2+)$  ions on the cytoplasmic side of the membrane and delivers them to the luminal side. The transfer of ions across the membrane is coupled to ATP hydrolysis and is associated with a transient phosphorylation that shifts the pump conformation from inward-facing to outward-facing state (PubMed:[16192278](#), PubMed:[16332677](#), PubMed:[30923126](#)). Plays a primary role in the maintenance of  $\text{Ca}(2+)$  homeostasis in the trans-Golgi compartment with a functional impact on Golgi and post-Golgi protein sorting as well as a structural impact on cisternae morphology (PubMed:[20439740](#), PubMed:[14632183](#)). Responsible for loading the Golgi stores with  $\text{Ca}(2+)$  ions in keratinocytes, contributing to keratinocyte differentiation and epidermis integrity (PubMed:[14632183](#), PubMed:[10615129](#), PubMed:[20439740](#)). Participates in  $\text{Ca}(2+)$  and  $\text{Mn}(2+)$  ions uptake into the Golgi store of hippocampal neurons and regulates protein trafficking required for neural polarity (By similarity). May also play a role in the maintenance of  $\text{Ca}(2+)$  and  $\text{Mn}(2+)$  homeostasis and signaling in the cytosol while preventing cytotoxicity (PubMed:[21187401](#)).

#### Cellular Location

Golgi apparatus, trans-Golgi network membrane; Multi-pass membrane protein. Golgi apparatus, Golgi stack membrane; Multi-pass membrane protein. Note=During neuron differentiation, shifts from juxtanuclear Golgi position to multiple Golgi structures distributed over the neural soma with a predominance in the apical dendritic trunk {ECO:0000250|UniProtKB:Q80XR2}

#### Tissue Location


Found in most tissues except colon, thymus, spleen and leukocytes (PubMed:15831496). Expressed in keratinocytes (at protein level) (PubMed:15831496, PubMed:14632183)

#### ATP2C1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

#### ATP2C1 Antibody (C-term) - Images



ATP2C1 Antibody (C-term) (Cat. #AP17155b) western blot analysis in T47D cell line lysates (35ug/lane). This demonstrates the ATP2C1 antibody detected the ATP2C1 protein (arrow).

#### **ATP2C1 Antibody (C-term) - Background**

The protein encoded by this gene belongs to the family of P-type cation transport ATPases. This magnesium-dependent enzyme catalyzes the hydrolysis of ATP coupled with the transport of the calcium. Defects in this gene cause Hailey-Hailey disease, an autosomal dominant disorder. Alternatively spliced transcript variants encoding different isoforms have been identified.

#### **ATP2C1 Antibody (C-term) - References**

Baron, S., et al. Biochim. Biophys. Acta 1798(8):1512-1521(2010)  
Davila, S., et al. Genes Immun. 11(3):232-238(2010)  
Tian, H., et al. J. Dermatol. Sci. 58(1):80-82(2010)  
Ding, Y.G., et al. Clin. Exp. Dermatol. 34 (8), E968-E971 (2009) :  
Nechama, M., et al. BMC Cell Biol. 10, 70 (2009) :