

UGT2B10 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17222B**Specification**

UGT2B10 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	P36537
Other Accession	O97951 , O75310 , O02663 , P16662 , NP_001066.1
Reactivity	Human
Predicted	Monkey
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	60774
Antigen Region	431-458

UGT2B10 Antibody (C-term) - Additional Information**Gene ID** 7365**Other Names**

UDP-glucuronosyltransferase 2B10, UDPGT 2B10, UGT2B10

Target/Specificity

This UGT2B10 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 431-458 amino acids from the C-terminal region of human UGT2B10.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

UGT2B10 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

UGT2B10 Antibody (C-term) - Protein Information**Name** UGT2B10

Function UDPGT is of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds.

Cellular Location


Microsome membrane; Single-pass membrane protein. Endoplasmic reticulum membrane; Single-pass membrane protein

UGT2B10 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

UGT2B10 Antibody (C-term) - Images



UGT2B10 Antibody (C-term) (Cat. #AP17222b) western blot analysis in 293 cell line lysates (35ug/lane). This demonstrates the UGT2B10 antibody detected the UGT2B10 protein (arrow).

UGT2B10 Antibody (C-term) - Background

UDPGT is of major importance in the conjugation and subsequent elimination of potentially toxic xenobiotics and endogenous compounds.

UGT2B10 Antibody (C-term) - References

- Chen, G., et al. Cancer Res. 70(19):7543-7552(2010)
Berg, J.Z., et al. Cancer Epidemiol. Biomarkers Prev. 19(6):1423-1431(2010)
Chen, G., et al. Pharmacogenet. Genomics 18(3):181-191(2008)
Chen, G., et al. Cancer Res. 67(19):9024-9029(2007)
Kaivosaari, S., et al. Mol. Pharmacol. 72(3):761-768(2007)