

MRPL2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17267B

Specification

MRPL2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q5T653
Other Accession	Q2TA12 , NP_057034.2
Reactivity	Human
Predicted	Bovine
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	33301
Antigen Region	236-264

MRPL2 Antibody (C-term) - Additional Information

Gene ID 51069

Other Names

39S ribosomal protein L2, mitochondrial, L2mt, MRP-L2, MRPL2

Target/Specificity

This MRPL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 236-264 amino acids from the C-terminal region of human MRPL2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MRPL2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.


MRPL2 Antibody (C-term) - Protein Information

Name MRPL2

Cellular Location
Mitochondrion**MRPL2 Antibody (C-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MRPL2 Antibody (C-term) - Images

MRPL2 Antibody (C-term) (Cat. #AP17267b) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the MRPL2 antibody detected the MRPL2 protein (arrow).

MRPL2 Antibody (C-term) - Background

Mammalian mitochondrial ribosomal proteins are encoded by nuclear genes and help in protein synthesis within the mitochondrion. Mitochondrial ribosomes (mitoribosomes) consist of a small 28S subunit and a large 39S subunit. They have an estimated 75% protein to rRNA composition compared to prokaryotic ribosomes, where this ratio is reversed. Another difference between mammalian mitoribosomes and prokaryotic ribosomes is that the latter contain a 5S rRNA. Among different species, the proteins comprising the mitoribosome differ greatly in sequence, and sometimes in biochemical properties, which prevents easy recognition by sequence homology. This gene encodes a 39S subunit protein that belongs to the EcoL2 ribosomal protein family. A pseudogene corresponding to this gene is found on chromosome 12q.

MRPL2 Antibody (C-term) - References

Lamesch, P., et al. Genomics 89(3):307-315(2007)

Mungall, A.J., et al. Nature 425(6960):805-811(2003)
Zhang, Z., et al. Genomics 81(5):468-480(2003)
Kenmochi, N., et al. Genomics 77 (1-2), 65-70 (2001) :
O'Brien, T.W., et al. J. Biol. Chem. 274(51):36043-36051(1999)