

Mouse Stk11 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17316b**Specification**

Mouse Stk11 Antibody (C-term) - Product Information

Application	IF, IHC-P, WB,E
Primary Accession	O9WTK7
Other Accession	NP_035622.1
Reactivity	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	49267
Antigen Region	393-420

Mouse Stk11 Antibody (C-term) - Additional Information**Gene ID** 20869**Other Names**Serine/threonine-protein kinase STK11, Liver kinase B1 homolog, LKB1, mLKB1, Stk11
{ECO:0000312|MGI:MGI:1341870}**Target/Specificity**

This Mouse Stk11 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 393-420 amino acids from the C-terminal region of mouse Stk11.

Dilution

IF~~1:25

IHC-P~~1:25

WB~~1:500-1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

Mouse Stk11 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

Mouse Stk11 Antibody (C-term) - Protein Information

Name Stk11 {ECO:0000312|MGI:MGI:1341870}

Function Tumor suppressor serine/threonine-protein kinase that controls the activity of AMP-activated protein kinase (AMPK) family members, thereby playing a role in various processes such as cell metabolism, cell polarity, apoptosis and DNA damage response. Acts by phosphorylating the T-loop of AMPK family proteins, thus promoting their activity: phosphorylates PRKAA1, PRKAA2, BRSK1, BRSK2, MARK1, MARK2, MARK3, MARK4, NUA1, NUA2, SIK1, SIK2, SIK3 and SNRK but not MELK. Also phosphorylates non-AMPK family proteins such as STRADA, PTEN and possibly p53/TP53. Acts as a key upstream regulator of AMPK by mediating phosphorylation and activation of AMPK catalytic subunits PRKAA1 and PRKAA2 and thereby regulates processes including: inhibition of signaling pathways that promote cell growth and proliferation when energy levels are low, glucose homeostasis in liver, activation of autophagy when cells undergo nutrient deprivation, and B-cell differentiation in the germinal center in response to DNA damage. Also acts as a regulator of cellular polarity by remodeling the actin cytoskeleton. Required for cortical neuron polarization by mediating phosphorylation and activation of BRSK1 and BRSK2, leading to axon initiation and specification. Involved in DNA damage response: interacts with p53/TP53 and recruited to the CDKN1A/WAF1 promoter to participate in transcription activation. Able to phosphorylate p53/TP53; the relevance of such result in vivo is however unclear and phosphorylation may be indirect and mediated by downstream STK11/LKB1 kinase NUA1. Also acts as a mediator of p53/TP53-dependent apoptosis via interaction with p53/TP53: translocates to the mitochondrion during apoptosis and regulates p53/TP53-dependent apoptosis pathways. Regulates UV radiation-induced DNA damage response mediated by CDKN1A. In association with NUA1, phosphorylates CDKN1A in response to UV radiation and contributes to its degradation which is necessary for optimal DNA repair (PubMed:[25329316](#)).

Cellular Location

Nucleus. Cytoplasm. Membrane. Mitochondrion. Note=Translocates to mitochondrion during apoptosis (By similarity). A small fraction localizes at membranes. Relocates to the cytoplasm when bound to STRAD (STRADA or STRADB) and CAB39/MO25 (CAB39/MO25alpha or CAB39L/MO25beta). PTEN promotes cytoplasmic localization (By similarity).

Tissue Location


[Isoform 1]: Widely expressed. [Isoform 3]: Expressed in adult brain and liver and absent from tissues derived from postnatal day 7


Mouse Stk11 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Mouse Stk11 Antibody (C-term) - Images







Fluorescent image of A549 cells stained with XAF1 Mouse Stk11 Antibody (C-term)(Cat#AP17316b). AP17316b was diluted at 1:100 dilution. An Alexa Fluor 488-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody (green). Cytoplasmic actin was counterstained with Alexa Fluor® 555 conjugated with Phalloidin (red).



Mouse Stk11 Antibody (C-term) (Cat. #AP17316b) western blot analysis in mouse cerebellum tissue lysates (35ug/lane). This demonstrates the Stk11 antibody detected the Stk11 protein (arrow).



Western blot analysis of lysate from NCI-H1299 cell line, using Mouse Stk11 Antibody (C-term)(Cat. #AP17316b). AP17316b was diluted at 1:1000. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug.



Immunohistochemical analysis of paraffin-embedded M.skeletal muscle section using Mouse Stk11 Antibody (C-term)(Cat#AP17316b). AP17316b was diluted at 1:25 dilution. A peroxidase-conjugated goat anti-rabbit IgG at 1:400 dilution was used as the secondary antibody, followed by DAB staining.

Mouse Stk11 Antibody (C-term) - Background

Essential role in G1 cell cycle arrest. Phosphorylates and activates members of the AMPK-related subfamily of protein kinases (By similarity). Tumor suppressor.

Mouse Stk11 Antibody (C-term) - References

Gao, Y., et al. Proc. Natl. Acad. Sci. U.S.A. 107(44):18892-18897(2010)
Bungard, D., et al. Science 329(5996):1201-1205(2010)
Koh, H.J., et al. Proc. Natl. Acad. Sci. U.S.A. 107(35):15541-15546(2010)
Foretz, M., et al. J. Clin. Invest. 120(7):2355-2369(2010)
Asada, N., et al. J. Neurosci. 30(26):8852-8865(2010)