

MLF1IP Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP17429b

Specification

MLF1IP Antibody (C-term) - Product Information

Application WB,E **Primary Accession** 071F23 Other Accession NP 078905.2 Reactivity Human Host **Rabbit** Clonality **Polyclonal** Isotype Rabbit IgG Calculated MW 47522 Antigen Region 292-319

MLF1IP Antibody (C-term) - Additional Information

Gene ID 79682

Other Names

Centromere protein U, CENP-U, Centromere protein of 50 kDa, CENP-50, Interphase centromere complex protein 24, KSHV latent nuclear antigen-interacting protein 1, MLF1-interacting protein, Polo-box-interacting protein 1, CENPU, ICEN24, KLIP1, MLF1IP, PBIP1

Target/Specificity

This MLF1IP antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 292-319 amino acids from the C-terminal region of human MLF1IP.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MLF1IP Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MLF1IP Antibody (C-term) - Protein Information

Name CENPU



Synonyms ICEN24, KLIP1, MLF1IP, PBIP1

Function Component of the CENPA-NAC (nucleosome-associated) complex, a complex that plays a central role in assembly of kinetochore proteins, mitotic progression and chromosome segregation. The CENPA-NAC complex recruits the CENPA-CAD (nucleosome distal) complex and may be involved in incorporation of newly synthesized CENPA into centromeres. Plays an important role in the correct PLK1 localization to the mitotic kinetochores. A scaffold protein responsible for the initial recruitment and maintenance of the kinetochore PLK1 population until its degradation. Involved in transcriptional repression.

Cellular Location

Cytoplasm. Nucleus. Chromosome, centromere, kinetochore. Note=Localizes in the kinetochore domain of centromeres Colocalizes with PLK1 at the interzone between the inner and the outer kinetochore plates

Tissue Location


Expressed at high levels in the testis, fetal liver, thymus, bone marrow and at lower levels in the lymph nodes, placenta, colon and spleen. Present in all cell lines examined, including B-cells, T-cells, epithelial cells and fibroblast cells Expressed at high levels in glioblastoma cell lines

MLF1IP Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

MLF1IP Antibody (C-term) - Images



MLF1IP Antibody (C-term) (Cat. #AP17429b) western blot analysis in WiDr cell line lysates (35ug/lane). This demonstrates the MLF1IP antibody detected the MLF1IP protein (arrow).

MLF1IP Antibody (C-term) - Background





The centromere is a specialized chromatin domain, present throughout the cell cycle, that acts as a platform on which the transient assembly of the kinetochore occurs during mitosis. All active centromeres are characterized by the presence of long arrays of nucleosomes in which CENPA (MIM 117139) replaces histone H3 (see MIM 601128). MLF1IP, or CENPU, is an additional factor required for centromere assembly (Foltz et al., 2006 [PubMed 16622419]).

MLF1IP Antibody (C-term) - References

Suzuki, H., et al. Anticancer Res. 27 (3B), 1423-1430 (2007): Kang, Y.H., et al. Mol. Cell 24(3):409-422(2006) lzuta, H., et al. Genes Cells 11(6):673-684(2006) Okada, M., et al. Nat. Cell Biol. 8(5):446-457(2006) Foltz, D.R., et al. Nat. Cell Biol. 8(5):458-469(2006)