

ZBTB22 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17466b**Specification**

ZBTB22 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O15209
Other Accession	O9Z0G7 , NP_001138810.1 , NP_005444.4
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	65602
Antigen Region	476-502

ZBTB22 Antibody (C-term) - Additional Information**Gene ID** 9278**Other Names**

Zinc finger and BTB domain-containing protein 22, Protein BING1, Zinc finger and BTB domain-containing protein 22A, Zinc finger protein 297, ZBTB22, BING1, ZBTB22A, ZNF297

Target/Specificity

This ZBTB22 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 476-502 amino acids from the C-terminal region of human ZBTB22.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZBTB22 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZBTB22 Antibody (C-term) - Protein Information**Name** ZBTB22

Synonyms BING1, ZBTB22A, ZNF297

Function May be involved in transcriptional regulation.


Cellular Location
Nucleus.

ZBTB22 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZBTB22 Antibody (C-term) - Images



Western blot analysis of lysates from MCF-7, SH-SY5Y cell line (from left to right), using ZBTB22 Antibody (C-term)(Cat. #AP17466b). AP17466b was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysates at 35ug per lane.

ZBTB22 Antibody (C-term) - Background

ZBTB22 may be involved in transcriptional regulation.

ZBTB22 Antibody (C-term) - References

Barcellos, L.F., et al. PLoS Genet. 5 (10), E1000696 (2009) :
Mungall, A.J., et al. Nature 425(6960):805-811(2003)
Herberg, J.A., et al. J. Mol. Biol. 277(4):839-857(1998)
Herberg, J.A., et al. Eur. J. Immunol. 28(2):459-467(1998)