

MTRF1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17473c

Specification

MTRF1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	O75570
Other Accession	O8K126 , Q3MHI7 , NP_004285.2
Reactivity	Human
Predicted	Bovine, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	52306
Antigen Region	232-259

MTRF1 Antibody (Center) - Additional Information

Gene ID 9617

Other Names

Peptide chain release factor 1, mitochondrial, MRF-1, MtRF-1, MTRF1

Target/Specificity

This MTRF1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 232-259 amino acids from the Central region of human MTRF1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MTRF1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

MTRF1 Antibody (Center) - Protein Information

Name RF1M

Function Mitochondrial peptide chain release factor that directs the termination of translation in

response to the peptide chain non- canonical stop codons AGG and AGA (PubMed:[36302763](#), PubMed:[36596788](#), PubMed:[37141370](#)). Non-canonical termination codons AGG and AGA are found at the end of MT-CO1/COX1 and MT-ND6/ND6 open reading frames, respectively (PubMed:[37141370](#)). Recognizes non-canonical stop codons via a network of interactions between the codon, MTRF1 and the ribosomal RNA (rRNA): in contrast to other translation release factors, which identify the codon in the A-site via direct interactions of amino acid side chains with the bases, MTRF1 repositions the first 2 bases of the stop codon to use an intricate network of interactions that includes residues of the release factor, the rRNA of the small ribosomal subunit, as well as neighboring bases of the mRNA (PubMed:[37141370](#)).

Cellular Location


Mitochondrion

MTRF1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MTRF1 Antibody (Center) - Images



Western blot analysis of lysate from human lung tissue lysate, using MTRF1 Antibody (Center)(Cat. #AP17473c). AP17473c was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.

MTRF1 Antibody (Center) - Background

The protein encoded by this gene was determined by in silico methods to be a mitochondrial protein with similarity to the peptide chain release factors (RFs) discovered in bacteria and

yeast. The peptide chain release factors direct the termination of translation in response to the peptide chain termination codons. Initially thought to have a role in the termination of mitochondria protein synthesis, a recent publication found no mitochondrial translation release functionality. Multiple alternatively spliced transcript variants have been suggested by mRNA and EST data; however, their full-length natures are not clear. [provided by RefSeq].

MTRF1 Antibody (Center) - References

Antonicka, H., et al. Am. J. Hum. Genet. 87(1):115-122(2010)
Nozaki, Y., et al. Genes Cells 13(5):429-438(2008)
Soleimanpour-Lichaei, H.R., et al. Mol. Cell 27(5):745-757(2007)
Hansen, L.L., et al. Cytogenet. Cell Genet. 88 (1-2), 91-92 (2000) :
Zhang, Y., et al. Biochim. Biophys. Acta 1443 (1-2), 245-250 (1998) :