

ATP6V1C2 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17823c**Specification**

ATP6V1C2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q8NEY4
Other Accession	NP_653184.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	48759
Antigen Region	276-304

ATP6V1C2 Antibody (Center) - Additional Information**Gene ID** 245973**Other Names**

V-type proton ATPase subunit C 2, V-ATPase subunit C 2, Vacuolar proton pump subunit C 2, ATP6V1C2

Target/Specificity

This ATP6V1C2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 276-304 amino acids from the Central region of human ATP6V1C2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ATP6V1C2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

ATP6V1C2 Antibody (Center) - Protein Information**Name** ATP6V1C2**Function** Subunit of the V1 complex of vacuolar(H⁺)-ATPase (V-ATPase), a multisubunit enzyme

composed of a peripheral complex (V1) that hydrolyzes ATP and a membrane integral complex (V0) that translocates protons (By similarity). V-ATPase is responsible for acidifying and maintaining the pH of intracellular compartments and in some cell types, is targeted to the plasma membrane, where it is responsible for acidifying the extracellular environment (By similarity). Subunit C is necessary for the assembly of the catalytic sector of the enzyme and is likely to have a specific function in its catalytic activity (By similarity).


Tissue Location

Kidney and placenta..

ATP6V1C2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ATP6V1C2 Antibody (Center) - Images

ATP6V1C2 Antibody (Center) (Cat. #AP17823c) western blot analysis in NCI-H292 cell line lysates (35ug/lane). This demonstrates the ATP6V1C2 antibody detected the ATP6V1C2 protein (arrow).

ATP6V1C2 Antibody (Center) - Background

This gene encodes a component of vacuolar ATPase (V-ATPase), a multisubunit enzyme that mediates acidification of eukaryotic intracellular organelles. V-ATPase dependent organelle acidification is necessary for such intracellular processes as protein sorting, zymogen activation, receptor-mediated endocytosis, and synaptic vesicle proton gradient generation. V-ATPase is composed of a cytosolic V1 domain and a transmembrane V0 domain. The V1 domain consists of three A, three B, and two G subunits, as well as a C, D, E, F, and H subunit. The V1 domain contains the ATP catalytic site. This gene encodes alternate transcriptional splice

variants, encoding different V1 domain C subunit isoforms.

ATP6V1C2 Antibody (Center) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Gruber, G. Biochem. Soc. Trans. 33 (PT 4), 883-885 (2005) :
Morel, N. Biol. Cell 95(7):453-457(2003)
Smith, A.N., et al. Mol. Cell 12(4):801-803(2003)
Kawasaki-Nishi, S., et al. FEBS Lett. 545(1):76-85(2003)