

WDR20 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17832a**Specification**

WDR20 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q8TBZ3
Other Accession	NP_653175.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	62893
Antigen Region	5-32

WDR20 Antibody (N-term) - Additional Information**Gene ID** 91833**Other Names**

WD repeat-containing protein 20, Protein DMR, WDR20

Target/Specificity

This WDR20 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 5-32 amino acids from the N-terminal region of human WDR20.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

WDR20 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

WDR20 Antibody (N-term) - Protein Information**Name** WDR20**Function** Regulator of deubiquitinating complexes. Activates deubiquitinating activity of

complexes containing USP12 (PubMed:[20147737](#), PubMed:[27373336](#)). Anchors at the base of the ubiquitin-contacting loop of USP12 and remotely modulates the catalytic center of the enzyme (PubMed:[27373336](#)). Regulates shuttling of the USP12 deubiquitinase complex between the plasma membrane, cytoplasm and nucleus (PubMed:[30466959](#)).

Cellular Location


Cytoplasm. Nucleus

WDR20 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

WDR20 Antibody (N-term) - Images



WDR20 Antibody (N-term) (Cat. #AP17832a) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the WDR20 antibody detected the WDR20 protein (arrow).

WDR20 Antibody (N-term) - Background

This gene encodes a WD repeat-containing protein. Multiple alternatively spliced transcript variants have been found for this gene, but the full-length nature of some variants has not been defined.

WDR20 Antibody (N-term) - References

Bailey, S.D., et al. Diabetes Care 33(10):2250-2253(2010)
Kee, Y., et al. J. Biol. Chem. 285(15):11252-11257(2010)
Talmud, P.J., et al. Am. J. Hum. Genet. 85(5):628-642(2009)
Olsen, J.V., et al. Cell 127(3):635-648(2006)