

OR2C3 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP17841b

Specification

OR2C3 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q8N628
Other Accession	NP_932340.3
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35376
Antigen Region	246-273

OR2C3 Antibody (C-term) - Additional Information

Gene ID 81472

Other Names

Olfactory receptor 2C3, Olfactory receptor 2C4, Olfactory receptor 2C5, Olfactory receptor OR1-30, OR2C3, OR2C4, OR2C5P

Target/Specificity

This OR2C3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 246-273 amino acids from the C-terminal region of human OR2C3.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR2C3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR2C3 Antibody (C-term) - Protein Information

Name OR2C3

Synonyms OR2C4, OR2C5P

Function Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR2C3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR2C3 Antibody (C-term) - Images



OR2C3 Antibody (C-term) (Cat. #AP17841b) western blot analysis in ZR-75-1 cell line lysates (35ug/lane). This demonstrates the OR2C3 antibody detected the OR2C3 protein (arrow).

OR2C3 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR2C3 Antibody (C-term) - References

Trevino, L.R., et al. Nat. Genet. 41(9):1001-1005(2009)

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)
Fuchs, T., et al. Genomics 80(3):295-302(2002)