

R4RL2 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18016B**Specification**

R4RL2 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q86UN3
Other Accession	NP_848665.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	46106
Antigen Region	279-308

R4RL2 Antibody (C-term) - Additional Information**Gene ID** 349667**Other Names**

Reticulon-4 receptor-like 2, Nogo receptor-like 3, Nogo-66 receptor homolog 1, Nogo-66 receptor-related protein 2, NgR2, RTN4RL2 (HGNC:23053)

Target/Specificity

This R4RL2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 279-308 amino acids from the C-terminal region of human R4RL2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

R4RL2 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

R4RL2 Antibody (C-term) - Protein Information**Name** RTN4RL2 ([HGNC:23053](#))

Function Cell surface receptor that plays a functionally redundant role in the inhibition of neurite outgrowth mediated by MAG (By similarity). Plays a functionally redundant role in postnatal brain development. Contributes to normal axon migration across the brain midline and normal formation of the corpus callosum. Does not seem to play a significant role in regulating axon regeneration in the adult central nervous system. Protects motoneurons against apoptosis; protection against apoptosis is probably mediated by MAG (By similarity). Like other family members, plays a role in restricting the number of dendritic spines and the number of synapses that are formed during brain development (PubMed:[22325200](#)). Signaling mediates activation of Rho and downstream reorganization of the actin cytoskeleton (PubMed:[22325200](#)).

Cellular Location

Cell membrane; Lipid-anchor, GPI-anchor. Membrane raft. Cell projection, dendrite {ECO:0000250|UniProtKB:Q7M6Z0}. Perikaryon {ECO:0000250|UniProtKB:Q80WD1}. Cell projection, axon {ECO:0000250|UniProtKB:Q80WD1}. Note=Localized to the surface of neurons, including axons. Detected close to synapses, but is excluded from synapses. {ECO:0000250|UniProtKB:Q7M6Z0}

Tissue Location


Highly expressed in brain and liver. Expressed at lower levels in kidney, mammary gland, placenta, skeletal muscle, spleen and thyroid.

R4RL2 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

R4RL2 Antibody (C-term) - Images



R4RL2 Antibody (C-term) (Cat. #AP18016b) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the R4RL2 antibody detected the R4RL2 protein (arrow).

R4RL2 Antibody (C-term) - Background

RTN4RL2 may play a role in regulating axonal regeneration and plasticity in the adult central nervous system.

R4RL2 Antibody (C-term) - References

Lee, H., et al. J. Neurosci. 28(11):2753-2765(2008)
Taylor, T.D., et al. Nature 440(7083):497-500(2006)
Walmsley, A.R., et al. Biochem. Biophys. Res. Commun. 327(1):112-116(2005)
Lauren, J., et al. Mol. Cell. Neurosci. 24(3):581-594(2003)
Barton, W.A., et al. EMBO J. 22(13):3291-3302(2003)