

### Mouse Stk24 Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18059c

### **Specification**

### Mouse Stk24 Antibody (Center) - Product Information

Application WB,E
Primary Accession Q99KH8

Other Accession Q9Y6E0, NP\_663440.1

Reactivity Human, Mouse

Host Rabbit
Clonality Polyclonal
Isotype Rabbit IgG
Calculated MW 47954
Antigen Region 222-249

### Mouse Stk24 Antibody (Center) - Additional Information

#### Gene ID 223255

### **Other Names**

Serine/threonine-protein kinase 24, Mammalian STE20-like protein kinase 3, MST-3, STE20-like kinase MST3, Serine/threonine-protein kinase 24 35 kDa subunit, Mammalian STE20-like protein kinase 3 N-terminal, MST3/N, Serine/threonine-protein kinase 24 12 kDa subunit, Mammalian STE20-like protein kinase 3 C-terminal, MST3/C, Stk24, Mst3, Stk3

### Target/Specificity

This Mouse Stk24 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 222-249 amino acids from the Central region of mouse Stk24.

# **Dilution**

WB~~1:1000

### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

### **Precautions**

Mouse Stk24 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

### Mouse Stk24 Antibody (Center) - Protein Information

Name Stk24



### Synonyms Mst3, Stk3

Function Serine/threonine-protein kinase that acts on both serine and threonine residues and promotes apoptosis in response to stress stimuli and caspase activation. Mediates oxidative-stress-induced cell death by modulating phosphorylation of JNK1-JNK2 (MAPK8 and MAPK9), p38 (MAPK11, MAPK12, MAPK13 and MAPK14) during oxidative stress. Plays a role in a staurosporine-induced caspase-independent apoptotic pathway by regulating the nuclear translocation of AIFM1 and ENDOG and the DNase activity associated with ENDOG. Phosphorylates STK38L on 'Thr-442' and stimulates its kinase activity. In association with STK26 negatively regulates Golgi reorientation in polarized cell migration upon RHO activation. Regulates also cellular migration with alteration of PTPN12 activity and PXN phosphorylation: phosphorylates PTPN12 and inhibits its activity and may regulate PXN phosphorylation through PTPN12. Acts as a key regulator of axon regeneration in the optic nerve and radial nerve (By similarity).

#### **Cellular Location**


Cytoplasm. Nucleus. Membrane. Note=The truncated form (MST3/N) translocates to the nucleus. Colocalizes with STK38L in the membrane (By similarity).

## Mouse Stk24 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

### Mouse Stk24 Antibody (Center) - Images



Mouse Stk24 Antibody (Center) (Cat. #AP18059c) western blot analysis in SK-BR-3 cell line lysates (35ug/lane). This demonstrates the Stk24 antibody detected the Stk24 protein (arrow).

### Mouse Stk24 Antibody (Center) - Background

Protein kinase that act on both serine and threonine residues (By similarity).



## Mouse Stk24 Antibody (Center) - References

Taniwaki, T., et al. Dev. Growth Differ. 47(3):163-172(2005) Shu, H., et al. Mol. Cell Proteomics 3(3):279-286(2004) Zambrowicz, B.P., et al. Proc. Natl. Acad. Sci. U.S.A. 100(24):14109-14114(2003) Piao, Y., et al. Genome Res. 11(9):1553-1558(2001) Araki, K., et al. Cell. Mol. Biol. (Noisy-le-grand) 45(5):737-750(1999)