

NPNT Antibody (Center)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18069c

Specification

NPNT Antibody (Center) - Product Information

Application WB,E
Primary Accession Q6UXI9

Other Accession NP_001028219.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Polyclonal
Rabbit IgG
61907
365-392

NPNT Antibody (Center) - Additional Information

Gene ID 255743

Other Names

Nephronectin, Preosteoblast EGF-like repeat protein with MAM domain, Protein EGFL6-like, NPNT, EGFL6L, POEM

Target/Specificity

This NPNT antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 365-392 amino acids from the Central region of human NPNT.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

NPNT Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

NPNT Antibody (Center) - Protein Information

Name NPNT



Tel: 858.875.1900 Fax: 858.875.1999

Synonyms EGFL6L, POEM

Function Functional ligand of integrin alpha-8/beta-1 in kidney development. Regulates the expression of GDNF with integrin alpha- 8/beta-1 which is essential for kidney development. May also play a role in the development and function of various tissues, regulating cell adhesion, spreading and survival through the binding of several integrins (By similarity).

Cellular Location

Secreted, extracellular space, extracellular matrix. Note=Trapped on the cell surface or in the extracellular matrix.

Tissue Location


Expressed in kidney and lung and to a lower extent in brain, pregnant uterus, placenta, thyroid gland and blood vessels

NPNT Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture


NPNT Antibody (Center) - Images



NPNT Antibody (Center) (Cat. #AP18069c) western blot analysis in MDA-MB453 cell line lysates (35ug/lane). This demonstrates the NPNT antibody detected the NPNT protein (arrow).

NPNT Antibody (Center) - Background

Functional ligand of integrin alpha-8/beta-1 in kidney development. Regulates the expression of GDNF with integrin alpha-8/beta-1 which is essential for kidney development. May also play a role in the development and function of various tissues, regulating cell adhesion, spreading and survival through the binding of several integrins (By similarity).





NPNT Antibody (Center) - References

Kalsi, G., et al. Hum. Mol. Genet. 19(12):2497-2506(2010) Hancock, D.B., et al. Nat. Genet. 42(1):45-52(2010) Kahai, S., et al. PLoS ONE 4 (10), E7535 (2009) : Huang, J.T., et al. Int. J. Mol. Med. 15(4):719-724(2005) Brandenberger, R., et al. J. Cell Biol. 154(2):447-458(2001)