

ONECUT3 Antibody (C-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18070b

Specification

ONECUT3 Antibody (C-term) - Product Information

Application WB,E
Primary Accession 060422

Other Accession <u>Q8K557</u>, <u>NP 001073957.1</u>

Reactivity
Predicted
Host
Clonality
Isotype
Calculated MW
Antigen Region

Human
Mouse
Rabbit
Polyclonal
Rabbit IgG
393-419

ONECUT3 Antibody (C-term) - Additional Information

Gene ID 390874

Other Names

One cut domain family member 3, One cut homeobox 3, Transcription factor ONECUT-3, OC-3, ONECUT3

Target/Specificity

This ONECUT3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 393-419 amino acids from the C-terminal region of human ONECUT3.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ONECUT3 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ONECUT3 Antibody (C-term) - Protein Information

Name ONECUT3



Function Transcriptional activator. Binds the consensus DNA sequence 5'-DHWATTGAYTWWD-3' on a variety of gene promoters such as those of HNF3B and TTR (By similarity).

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000255|PROSITE-ProRule:PRU00374}

ONECUT3 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

ONECUT3 Antibody (C-term) - Images



Anti-ONECUT3 Antibody (C-term) at 1:1000 dilution + MDA-MB-231 whole cell lysate Lysates/proteins at 20 μ g per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 50 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

ONECUT3 Antibody (C-term) - Background

ONECUT3 is transcriptional activator. It binds the consensus DNA sequence 5'-DHWATTGAYTWWD-3' on a variety of gene promoters such as those of HNF3B and TTR (By similarity).

ONECUT3 Antibody (C-term) - References

Grimwood, J., et al. Nature 428(6982):529-535(2004) Vanhorenbeeck, V., et al. Biochem. Biophys. Res. Commun. 292(4):848-854(2002) Venter, J.C., et al. Science 291(5507):1304-1351(2001) Jacquemin, P., et al. J. Biol. Chem. 274(5):2665-2671(1999)