

ZNF187 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18078a**Specification**

ZNF187 Antibody (N-term) - Product Information

Application	IF, WB,E
Primary Accession	Q16670
Other Accession	NP_001018854.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	55254
Antigen Region	3-31

ZNF187 Antibody (N-term) - Additional Information**Gene ID** 7741**Other Names**

Zinc finger and SCAN domain-containing protein 26, Protein SRE-ZBP, Zinc finger protein 187, ZSCAN26, ZNF187

Target/Specificity

This ZNF187 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 3-31 amino acids from the N-terminal region of human ZNF187.

Dilution

IF~~1:10~50

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

ZNF187 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ZNF187 Antibody (N-term) - Protein Information**Name** ZSCAN26

Synonyms ZNF187

Function May be involved in transcriptional regulation.


Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187, ECO:0000269|PubMed:1569959}


ZNF187 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF187 Antibody (N-term) - Images

Fluorescent image of A549 cell stained with ZNF187 Antibody (N-term)(Cat#AP18078a).A549 cells were fixed with 4% PFA (20 min), permeabilized with Triton X-100 (0.1%, 10 min), then incubated with ZNF187 primary antibody (1:25, 1 h at 37°C). For secondary antibody, Alexa Fluor® 488 conjugated donkey anti-rabbit antibody (green) was used (1:400, 50 min at 37°C).Cytoplasmic actin was counterstained with Alexa Fluor® 555 (red) conjugated Phalloidin (7units/ml, 1 h at 37°C).ZNF187 immunoreactivity is localized to Nucleus significantly.



ZNF187 Antibody (N-term) (Cat. #AP18078a) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the ZNF187 antibody detected the ZNF187 protein (arrow).

ZNF187 Antibody (N-term) - Background

ZNF187 may be involved in transcriptional regulation (Probable).

ZNF187 Antibody (N-term) - References

Attar, R.M., et al. Mol. Cell. Biol. 12(5):2432-2443(1992)