

OR10AD1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18196c**Specification**

OR10AD1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q8NGE0
Other Accession	NP_001004134.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35698
Antigen Region	149-177

OR10AD1 Antibody (Center) - Additional Information**Gene ID** 121275**Other Names**

Olfactory receptor 10AD1, Olfactory receptor OR12-1, OR10AD1, OR10AD1P

Target/Specificity

This OR10AD1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 149-177 amino acids from the Central region of human OR10AD1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR10AD1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

OR10AD1 Antibody (Center) - Protein Information**Name** OR10AD1**Synonyms** OR10AD1P

Function Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR10AD1 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR10AD1 Antibody (Center) - Images



OR10AD1 Antibody (Center) (Cat. #AP18196c) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the OR10AD1 antibody detected the OR10AD1 protein (arrow).

OR10AD1 Antibody (Center) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR10AD1 Antibody (Center) - References

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)

Fuchs, T., et al. Genomics 80(3):295-302(2002)