

MED18 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18275B

Specification

MED18 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	Q9BUE0
Other Accession	Q9CZ82 , Q7T3H7 , Q0VCD4 , NP_001120822.1
Reactivity	Human
Predicted	Bovine, Zebrafish, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	23663
Antigen Region	180-206

MED18 Antibody (C-term) - Additional Information

Gene ID 54797

Other Names

Mediator of RNA polymerase II transcription subunit 18, Mediator complex subunit 18, p28b, MED18

Target/Specificity

This MED18 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 180-206 amino acids from the C-terminal region of human MED18.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

MED18 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

MED18 Antibody (C-term) - Protein Information

Name MED18

Function Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene- specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Cellular Location


Nucleus.

MED18 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

MED18 Antibody (C-term) - Images



MED18 Antibody (C-term) (Cat. #AP18275b) western blot analysis in HL-60 cell line lysates (35ug/lane). This demonstrates the MED18 antibody detected the MED18 protein (arrow).

MED18 Antibody (C-term) - Background

MED18 is a component of the Mediator complex, which is a coactivator for DNA-binding factors that activate transcription via RNA polymerase II (Sato et al., 2003 [PubMed 12584197]).[supplied by OMIM].

MED18 Antibody (C-term) - References

- Sato, S., et al. Mol. Cell 14(5):685-691(2004)
Tomomori-Sato, C., et al. J. Biol. Chem. 279(7):5846-5851(2004)
Sato, S., et al. J. Biol. Chem. 278(50):49671-49674(2003)

Sato, S., et al. J. Biol. Chem. 278(17):15123-15127(2003)