

#### **ZFP57 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18299C

## **Specification**

## **ZFP57 Antibody (Center) - Product Information**

Application WB,E
Primary Accession O9NU63

Other Accession NP 001103279.2

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region

Human
Rabbit
Rabbit
Rabbit
Polyclonal
Rabbit IgG
269-296

## **ZFP57 Antibody (Center) - Additional Information**

#### Gene ID 346171

#### **Other Names**

Zinc finger protein 57 homolog, Zfp-57, Zinc finger protein 698, ZFP57, C6orf40, ZNF698

#### Target/Specificity

This ZFP57 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 269-296 amino acids from the Central region of human ZFP57.

## **Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

#### **Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

## **Precautions**

ZFP57 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

## **ZFP57 Antibody (Center) - Protein Information**

Name ZFP57 (HGNC:18791)

Synonyms C6orf40, ZNF698



**Function** Transcription regulator required to maintain maternal and paternal gene imprinting, a process by which gene expression is restricted in a parent of origin-specific manner by epigenetic modification of genomic DNA and chromatin, including DNA methylation. Acts by controlling DNA methylation during the earliest multicellular stages of development at multiple imprinting control regions (ICRs) (PubMed:18622393, PubMed:30602440). Acts together with ZNF445, but ZNF445 seems to be the major factor in human early embryonic imprinting maintenance. In contrast, in mice, ZFP57 plays the predominant role in imprinting maintenance (PubMed:30602440). Required for the establishment of maternal methylation imprints at SNRPN locus. Acts as a transcriptional repressor in Schwann cells. Binds to a 5'-TGCCGC-3' consensus sequence and recognizes the methylated CpG within this element (By similarity).

#### **Cellular Location**


Nucleus. Note=Binds various differentially methylated regions (DMR)

## **ZFP57 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture


## ZFP57 Antibody (Center) - Images



Anti-ZFP57 Antibody (Center) at 1:1000 dilution + human placenta lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 52 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

## ZFP57 Antibody (Center) - Background

The protein encoded by this gene is a zinc finger protein containing a KRAB domain. Studies in mouse suggest that this





protein may function as a transcriptional repressor. Mutations in this gene have been associated with transient neonatal diabetes mellitus type 1 (TNDM1).

# **ZFP57 Antibody (Center) - References**

Spengler, S., et al. Eur J Med Genet 52(6):415-416(2009) Barcellos, L.F., et al. PLoS Genet. 5 (10), E1000696 (2009): Tse, K.P., et al. Am. J. Hum. Genet. 85(2):194-203(2009) Li, X., et al. Dev. Cell 15(4):547-557(2008) Mackay, D.J., et al. Nat. Genet. 40(8):949-951(2008)