

OR2AG2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18307a**Specification**

OR2AG2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	A6NM03
Other Accession	O9H205 , NP_001004490.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	35270
Antigen Region	64-90

OR2AG2 Antibody (N-term) - Additional Information**Gene ID** 338755**Other Names**

Olfactory receptor 2AG2, OR2AG2, OR2AG2P

Target/Specificity

This OR2AG2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 64-90 amino acids from the N-terminal region of human OR2AG2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR2AG2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR2AG2 Antibody (N-term) - Protein Information**Name** OR2AG2**Synonyms** OR2AG2P

Function Odorant receptor.

Cellular Location


Cell membrane; Multi-pass membrane protein.

OR2AG2 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR2AG2 Antibody (N-term) - Images



OR2AG2 Antibody (N-term) (Cat. #AP18307a) western blot analysis in K562 cell line lysates (35ug/lane). This demonstrates the OR2AG2 antibody detected the OR2AG2 protein (arrow).

OR2AG2 Antibody (N-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR2AG2 Antibody (N-term) - References

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)

Fuchs, T., et al. Genomics 80(3):295-302(2002)