

GNRHR Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18314a

Specification

GNRHR Antibody (N-term) - Product Information

WB,E
<u>P30968</u>
<u>NP_000397</u>
Human
Rabbit
Polyclonal
Rabbit IgG
37731
49-77

GNRHR Antibody (N-term) - Additional Information

Gene ID 2798

Other Names Gonadotropin-releasing hormone receptor, GnRH receptor, GnRH-R, GNRHR, GRHR

Target/Specificity

This GNRHR antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 49-77 amino acids from the N-terminal region of human GNRHR.

Dilution WB~~1:1000 E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

GNRHR Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

GNRHR Antibody (N-term) - Protein Information

Name GNRHR

Synonyms GRHR



Function Receptor for gonadotropin releasing hormone (GnRH) that mediates the action of GnRH to stimulate the secretion of the gonadotropic hormones luteinizing hormone (LH) and follicle-stimulating hormone (FSH). This receptor mediates its action by association with G-proteins that activate a phosphatidylinositol-calcium second messenger system. Isoform 2 may act as an inhibitor of GnRH-R signaling.

Cellular Location Cell membrane; Multi-pass membrane protein.

Tissue Location Pituitary, ovary, testis, breast and prostate but not in liver and spleen

GNRHR Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

GNRHR Antibody (N-term) - Images



Western blot analysis of lysate from human ovary tissue lysate, using GNRHR Antibody (N-term)(Cat. #AP18314a). AP18314a was diluted at 1:1000 at each lane. A goat anti-rabbit IgG H&L(HRP) at 1:5000 dilution was used as the secondary antibody. Lysate at 35ug per lane.

GNRHR Antibody (N-term) - Background

This gene encodes the receptor for type 1 gonadotropin-releasing hormone. This receptor is a member of the seven-transmembrane, G-protein coupled receptor (GPCR) family. It is expressed on the surface of pituitary gonadotrope cells as well



as lymphocytes, breast, ovary, and prostate. Following binding of gonadotropin-releasing hormone, the receptor associates with G-proteins that activate a phosphatidylinositol-calcium second messenger system. Activation of the receptor ultimately causes the release of gonadotropic luteinizing hormone (LH) and follicle stimulating hormone (FSH). Defects in this gene are a cause of hypogonadotropic hypogonadism (HH). Alternative splicing results in multiple transcript variants encoding different isoforms. More than 18 transcription initiation sites in the 5' region and multiple polyA signals in the 3' region have been identified for this gene.

GNRHR Antibody (N-term) - References

Canzian, F., et al. Hum. Mol. Genet. 19(19):3873-3884(2010) Armstrong, S.P., et al. J. Biol. Chem. 285(32):24360-24371(2010) Lee, G., et al. Cancer Immunol. Immunother. 59(7):1011-1019(2010) Re, M., et al. PLoS ONE 5 (7), E11489 (2010) : Armstrong, S.P., et al. J. Biol. Chem. 284(51):35746-35757(2009)