

**OR51G1 Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP18334b**

### Specification

#### OR51G1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	<a href="#">Q8NGK1</a>
Other Accession	<a href="#">NP_001005237.1</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36290
Antigen Region	287-316

#### OR51G1 Antibody (C-term) - Additional Information

##### Gene ID 79324

##### Other Names

Olfactory receptor 51G1, Olfactory receptor 51G3, Olfactory receptor OR11-29, OR51G1, OR51G3P

##### Target/Specificity

This OR51G1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 287-316 amino acids from the C-terminal region of human OR51G1.

##### Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

##### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

##### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

##### Precautions

OR51G1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

#### OR51G1 Antibody (C-term) - Protein Information

**Name** OR51G1

**Synonyms** OR51G3P

**Function** Odorant receptor.

**Cellular Location**


Cell membrane; Multi-pass membrane protein.

### OR51G1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### OR51G1 Antibody (C-term) - Images



OR51G1 Antibody (C-term) (Cat. #AP18334b) western blot analysis in ZR-75-1 cell line lysates (35ug/lane). This demonstrates the OR51G1 Antibody detected the OR51G1 protein (arrow).

### OR51G1 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

### OR51G1 Antibody (C-term) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)