

SCML1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18386a

Specification

SCML1 Antibody (N-term) - Product Information

Application WB,E
Primary Accession Q9UN30

Other Accession NP_001032624.1

Reactivity
Host
Clonality
Polyclonal
Isotype
Calculated MW
Antigen Region
Human
Rabbit
Rabbit
Rabbit
Polyclonal
Rabbit IgG
37447
Antigen Region

SCML1 Antibody (N-term) - Additional Information

Gene ID 6322

Other Names

Sex comb on midleg-like protein 1, SCML1

Target/Specificity

This SCML1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 15-41 amino acids from the N-terminal region of human SCML1.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

SCML1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

SCML1 Antibody (N-term) - Protein Information

Name SCML1

Function Putative Polycomb group (PcG) protein. PcG proteins act by forming multiprotein complexes, which are required to maintain the transcriptionally repressive state of homeotic



genes throughout development. May be involved in spermatogenesis during sexual maturation (By similarity).

Cellular Location Nucleus.

Tissue Location


Ubiquitous. Expressed in fetal and adult tissues.

SCML1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

SCML1 Antibody (N-term) - Images



SCML1 Antibody (N-term) (Cat. #AP18386a) western blot analysis in MCF-7 cell line lysates (35ug/lane). This demonstrates the SCML1 Antibody detected the SCML1 protein (arrow).

SCML1 Antibody (N-term) - Background

Putative Polycomb group (PcG) protein. PcG proteins act by forming multiprotein complexes, which are required to maintain the transcriptionally repressive state of homeotic genes throughout development. May be involved in spermatogenesis during sexual maturation (By similarity).

SCML1 Antibody (N-term) - References

Wu, H.H., et al. BMC Evol. Biol. 8, 192 (2008): Ross, M.T., et al. Nature 434(7031):325-337(2005) van de Vosse, E., et al. Genomics 49(1):96-102(1998)