

OR2W1 Antibody (C-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18506b**Specification**

OR2W1 Antibody (C-term) - Product Information

Application	WB,E
Primary Accession	O9Y3N9
Other Accession	NP_112165.1
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	36101
Antigen Region	245-272

OR2W1 Antibody (C-term) - Additional Information**Gene ID** 26692**Other Names**

Olfactory receptor 2W1, Hs6M1-15, Olfactory receptor OR6-13, OR2W1

Target/Specificity

This OR2W1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 245-272 amino acids from the C-terminal region of human OR2W1.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

OR2W1 Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

OR2W1 Antibody (C-term) - Protein Information**Name** OR2W1**Function** Odorant receptor.


Cellular Location

Cell membrane; Multi-pass membrane protein.

OR2W1 Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

OR2W1 Antibody (C-term) - Images

OR2W1 Antibody (C-term) (Cat. #AP18506b) western blot analysis in A549 cell line lysates (35ug/lane). This demonstrates the OR2W1 antibody detected the OR2W1 protein (arrow).

OR2W1 Antibody (C-term) - Background

Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptor proteins are members of a large family of G-protein-coupled receptors (GPCR) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone receptors and are responsible for the recognition and G protein-mediated transduction of odorant signals. The olfactory receptor gene family is the largest in the genome. The nomenclature assigned to the olfactory receptor genes and proteins for this organism is independent of other organisms.

OR2W1 Antibody (C-term) - References

Barcellos, L.F., et al. PLoS Genet. 5 (10), E1000696 (2009) :

Malnic, B., et al. Proc. Natl. Acad. Sci. U.S.A. 101(8):2584-2589(2004)
Mungall, A.J., et al. Nature 425(6960):805-811(2003)
Fuchs, T., et al. Genomics 80(3):295-302(2002)