

RAB40C Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18572c

Specification

RAB40C Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q96S21
Other Accession	Q8VHO4 , NP_001166134.1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	31304
Antigen Region	122-148

RAB40C Antibody (Center) - Additional Information

Gene ID 57799

Other Names

Ras-related protein Rab-40C, Rar-like protein, Ras-like protein family member 8C, SOCS box-containing protein RAR3, RAB40C, RARL, RASL8C

Target/Specificity

This RAB40C antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 122-148 amino acids from the Central region of human RAB40C.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

RAB40C Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

RAB40C Antibody (Center) - Protein Information

Name RAB40C ([HGNC:18285](#))

Synonyms RARL, RASL8C

Function RAB40C small GTPase acts as substrate-recognition component of the ECS(RAB40C) E3 ubiquitin ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins (PubMed:[15601820](#), PubMed:[35512830](#)). The Rab40 subfamily belongs to the Rab family that are key regulators of intracellular membrane trafficking, from the formation of transport vesicles to their fusion with membranes. Rabs cycle between an inactive GDP-bound form and an active GTP-bound form that is able to recruit to membranes different sets of downstream effectors directly responsible for vesicle formation, movement, tethering and fusion (PubMed:[29156729](#)). As part of the ECS(RAB40C) complex, mediates ANKRD28 ubiquitination and degradation, thereby inhibiting protein phosphatase 6 (PP6) complex activity and focal adhesion assembly during cell migration (PubMed:[35512830](#)). Also negatively regulate lipid droplets accumulation in a GTP-dependent manner (PubMed:[29156729](#)).

Cellular Location


Cell membrane; Lipid-anchor; Cytoplasmic side. Cytoplasm, cytosol. Golgi apparatus membrane. Note=Mostly localized in the cytosol and also with actin ruffles.

RAB40C Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

RAB40C Antibody (Center) - Images



RAB40C Antibody (Center) (Cat. #AP18572c) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the RAB40C antibody detected the RAB40C protein (arrow).



RAB40C Antibody (Center) (Cat. #AP18572c) western blot analysis in mouse bladder tissue lysates (35ug/lane). This demonstrates the RAB40C antibody detected the RAB40C protein (arrow).

RAB40C Antibody (Center) - Background

Probable substrate-recognition component of a SCF-like ECS (Elongin-Cullin-SOCS-box protein) E3 ubiquitin ligase complex which mediates the ubiquitination and subsequent proteasomal degradation of target proteins.

RAB40C Antibody (Center) - References

- Ng, E.L., et al. Brain Res Rev 58(1):236-246(2008)
- Lettre, G., et al. Nat. Genet. 40(5):584-591(2008)
- Martin, J., et al. Nature 432(7020):988-994(2004)
- Kile, B.T., et al. Trends Biochem. Sci. 27(5):235-241(2002)
- Pereira-Leal, J.B., et al. J. Mol. Biol. 313(4):889-901(2001)