

**H2BFS Antibody (Center)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP18574c****Specification**

---

**H2BFS Antibody (Center) - Product Information**

Application	WB,E
Primary Accession	<a href="#">P57053</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	13944
Antigen Region	58-84

**H2BFS Antibody (Center) - Additional Information****Gene ID** 102724334**Other Names**

Histone H2B type F-S, Histone H2Bs, H2B/s, H2BFS

**Target/Specificity**

This H2BFS antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 58-84 amino acids from the Central region of human H2BFS.

**Dilution**

WB~~1:1000

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

H2BFS Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

**H2BFS Antibody (Center) - Protein Information****Name** H2BC12L ([HGNC:4762](#))

**Function** Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and

chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

#### **Cellular Location**


Nucleus. Chromosome.

### **H2BFS Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **H2BFS Antibody (Center) - Images**



All lanes : Anti-H2BFS Antibody (Center) at 1:1000 dilution Lane 1: A431 whole cell lysate Lane 2: WiDr whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 14 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

### **H2BFS Antibody (Center) - Background**

Core component of nucleosome. Nucleosomes wrap and compact DNA into chromatin, limiting DNA accessibility to the cellular machineries which require DNA as a template. Histones thereby play a central role in transcription regulation, DNA repair, DNA replication and chromosomal stability. DNA accessibility is regulated via a complex set of post-translational modifications of histones, also called histone code, and nucleosome remodeling.

Has broad antibacterial activity. May contribute to the formation of the functional antimicrobial barrier of the colonic epithelium, and to the bactericidal activity of amniotic fluid.