

BRD9 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18588A**Specification**

BRD9 Antibody (N-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	O9H8M2
Other Accession	NP_001009877.2
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	3-29

BRD9 Antibody (N-term) - Additional Information**Gene ID** 65980**Other Names**

Bromodomain-containing protein 9, Rhabdomyosarcoma antigen MU-RMS-408, BRD9

Target/Specificity

This BRD9 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 3-29 amino acids from the N-terminal region of human BRD9.

DilutionWB~~1:1000
IHC-P~~1:100**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

BRD9 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

BRD9 Antibody (N-term) - Protein Information**Name** BRD9

Function Plays a role in chromatin remodeling and regulation of transcription (PubMed:[22464331](#), PubMed:[26365797](#)). Acts as a chromatin reader that recognizes and binds acylated histones: binds

histones that are acetylated and/or butyrylated (PubMed:[26365797](#)). Component of SWI/SNF chromatin remodeling subcomplex GBAF that carries out key enzymatic activities, changing chromatin structure by altering DNA- histone contacts within a nucleosome in an ATP-dependent manner (PubMed:[29374058](#)). Orchestrates also the RAD51-RAD54 complex formation and thereby plays a role in homologous recombination (HR) (PubMed:[32457312](#)).

Cellular Location


Nucleus.

BRD9 Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


BRD9 Antibody (N-term) - Images




All lanes : Anti-BRD9 Antibody (N-term) at 1:1000-1:2000 dilution Lane 1: HeLa whole cell lysate Lane 2: SH-SY5Y whole cell lysate Lane 3: Human placenta lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 67 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



BRD9 Antibody (N-term) (Cat. #AP18588a) western blot analysis in mouse Neuro-2a cell line lysates (35ug/lane). This demonstrates the BRD9 antibody detected the BRD9 protein (arrow).



Immunohistochemical analysis of AP18588A on paraffin-embedded Human placenta tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9.0). Samples were incubated with primary antibody (1:100) for 1 hour at room temperature. Undiluted CRP Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.



Immunohistochemical analysis of AP18588A on paraffin-embedded Human testis tissue. Tissue was fixed with formaldehyde at room temperature. Heat induced epitope retrieval was performed by EDTA buffer (pH9. 0). Samples were incubated with primary antibody(1:100) for 1 hour at room temperature. Undiluted CRF Anti-Polyvalent HRP Polymer antibody was used as the secondary antibody.

BRD9 Antibody (N-term) - Background

BRD9 is a bromodomain containing protein, which are known to bind to acetylated lysine residues.

BRD9 Antibody (N-term) - References

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) :
Scotto, L., et al. Mol. Cancer 7, 58 (2008) :
Clark, H.F., et al. Genome Res. 13(10):2265-2270(2003)