

### **UHRF2 Antibody (Center)**

Affinity Purified Rabbit Polyclonal Antibody (Pab) Catalog # AP18663c

### Specification

# **UHRF2 Antibody (Center) - Product Information**

Application
Primary Accession
Other Accession
Reactivity
Host
Clonality
Isotype
Calculated MW
Antigen Region

WB,E <u>Q96PU4</u> <u>NP\_690856.1</u> Human Rabbit Polyclonal Rabbit IgG 89985 479-508

# **UHRF2** Antibody (Center) - Additional Information

### Gene ID 115426

### **Other Names**

E3 ubiquitin-protein ligase UHRF2, 632-, Np95/ICBP90-like RING finger protein, Np95-like RING finger protein, Nuclear protein 97, Nuclear zinc finger protein Np97, RING finger protein 107, Ubiquitin-like PHD and RING finger domain-containing protein 2, Ubiquitin-like-containing PHD and RING finger domains protein 2, UHRF2, NIRF, RNF107

### Target/Specificity

This UHRF2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 479-508 amino acids from the Central region of human UHRF2.

**Dilution** WB~~1:1000 E~~Use at an assay dependent concentration.

#### Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

#### Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

#### **Precautions**

UHRF2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

# **UHRF2 Antibody (Center) - Protein Information**



## Name UHRF2

Synonyms NIRF, RNF107

**Function** E3 ubiquitin ligase that plays important roles in DNA methylation, histone modifications, cell cycle and DNA repair (PubMed:<u>15178429</u>, PubMed:<u>23404503</u>, PubMed:<u>27743347</u>, PubMed:<u>29506131</u>). Acts as a specific reader for 5-hydroxymethylcytosine (5hmC) and thereby recruits various substrates to these sites to ubiquitinate them (PubMed:<u>24813944</u>, PubMed:<u>27129234</u>). This activity also allows the maintenance of 5mC levels at specific genomic loci and regulates neuron-related gene expression (By similarity). Participates in cell cycle regulation by ubiquitinating cyclins CCND1 and CCNE1 and thereby inducing G1 arrest (PubMed:<u>15178429</u>, PubMed:<u>15361834</u>, PubMed:<u>21952639</u>). Also ubiquitinates PCNP leading to its degradation by the proteasome (PubMed:<u>12176013</u>, PubMed:<u>14741369</u>). Plays an active role in DNA damage repair by ubiquitinating p21/CDKN1A leading to its proteasomal degradation (PubMed:<u>29923055</u>). Also promotes DNA repair by acting as an interstrand cross-links (ICLs) sensor. Mechanistically, cooperates with UHRF1 to ensure recruitment of FANCD2 to ICLs, leading to FANCD2 monoubiquitination and subsequent activation (PubMed:<u>30335751</u>). Contributes to UV-induced DNA damage response by physically interacting with ATR in response to irradiation, thereby promoting ATR activation (PubMed:<u>33848395</u>).

### **Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00358, ECO:0000269|PubMed:12176013, ECO:0000269|PubMed:23404503, ECO:0000269|PubMed:27129234, ECO:0000269|PubMed:27743347, ECO:0000269|PubMed:29923055, ECO:0000269|PubMed:30335751}. Chromosome. Note=Enriched at genomic loci that are enriched for 5-hydroxymethylcytosine (5hmC)


# **UHRF2 Antibody (Center) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

UHRF2 Antibody (Center) - Images





UHRF2 Antibody (Center) (Cat. #AP18663c) western blot analysis in Jurkat cell line lysates (35ug/lane).This demonstrates the UHRF2 antibody detected the UHRF2 protein (arrow).

# UHRF2 Antibody (Center) - Background

This gene encodes a nuclear protein which is involved in cell-cycle regulation. The encoded protein is a ubiquitin-ligase capable of ubiquinating PCNP (PEST-containing nuclear protein), and together they may play a role in tumorigenesis. [provided by RefSeq].

## **UHRF2 Antibody (Center) - References**

Rose, J.E., et al. Mol. Med. 16 (7-8), 247-253 (2010) : He, X., et al. FEBS Lett. 583(21):3501-3507(2009) Iwata, A., et al. J. Biol. Chem. 284(15):9796-9803(2009) Unoki, M., et al. Oncogene 23(46):7601-7610(2004) Colland, F., et al. Genome Res. 14(7):1324-1332(2004)