

FOXP3 Antibody (C-term S418)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18718b**Specification**

FOXP3 Antibody (C-term S418) - Product Information

| | |
|-------------------|--|
| Application | WB,E |
| Primary Accession | O9BZS1 |
| Other Accession | O6U8D7 , NP_054728 |
| Reactivity | Human |
| Predicted | Monkey |
| Host | Rabbit |
| Clonality | Polyclonal |
| Isotype | Rabbit IgG |
| Calculated MW | 47244 |
| Antigen Region | 397-423 |

FOXP3 Antibody (C-term S418) - Additional Information**Gene ID** 50943**Other Names**

Forkhead box protein P3, Scurfin, FOXP3, IPEX

Target/Specificity

This FOXP3 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 397-423 amino acids from the C-terminal region of human FOXP3.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

FOXP3 Antibody (C-term S418) is for research use only and not for use in diagnostic or therapeutic procedures.

FOXP3 Antibody (C-term S418) - Protein Information**Name** FOXP3

Synonyms IPEX

Function Transcriptional regulator which is crucial for the development and inhibitory function of regulatory T-cells (Treg) (PubMed:[17377532](#), PubMed:[21458306](#), PubMed:[23947341](#), PubMed:[24354325](#), PubMed:[24722479](#), PubMed:[24835996](#), PubMed:[30513302](#), PubMed:[32644293](#)). Plays an essential role in maintaining homeostasis of the immune system by allowing the acquisition of full suppressive function and stability of the Treg lineage, and by directly modulating the expansion and function of conventional T-cells (PubMed:[23169781](#)). Can act either as a transcriptional repressor or a transcriptional activator depending on its interactions with other transcription factors, histone acetylases and deacetylases (PubMed:[17377532](#), PubMed:[21458306](#), PubMed:[23947341](#), PubMed:[24354325](#), PubMed:[24722479](#)). The suppressive activity of Treg involves the coordinate activation of many genes, including CTLA4 and TNFRSF18 by FOXP3 along with repression of genes encoding cytokines such as interleukin-2 (IL2) and interferon-gamma (IFNG) (PubMed:[17377532](#), PubMed:[21458306](#), PubMed:[23947341](#), PubMed:[24354325](#), PubMed:[24722479](#)). Inhibits cytokine production and T-cell effector function by repressing the activity of two key transcription factors, RELA and NFATC2 (PubMed:[15790681](#)). Mediates transcriptional repression of IL2 via its association with histone acetylase KAT5 and histone deacetylase HDAC7 (PubMed:[17360565](#)). Can activate the expression of TNFRSF18, IL2RA and CTLA4 and repress the expression of IL2 and IFNG via its association with transcription factor RUNX1 (PubMed:[17377532](#)). Inhibits the differentiation of IL17 producing helper T-cells (Th17) by antagonizing RORC function, leading to down-regulation of IL17 expression, favoring Treg development (PubMed:[18368049](#)). Inhibits the transcriptional activator activity of RORA (PubMed:[18354202](#)). Can repress the expression of IL2 and IFNG via its association with transcription factor IKZF4 (By similarity).

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00089, ECO:0000269|PubMed:17360565, ECO:0000269|PubMed:18354202, ECO:0000269|PubMed:22678915, ECO:0000269|PubMed:23396208, ECO:0000269|PubMed:23973222, ECO:0000269|PubMed:23973223, ECO:0000269|PubMed:32644293}. Cytoplasm Note=Predominantly expressed in the cytoplasm in activated conventional T-cells whereas predominantly expressed in the nucleus in regulatory T- cells (Treg). The 41 kDa form derived by proteolytic processing is found exclusively in the chromatin fraction of activated Treg cells (By similarity). {ECO:0000250|UniProtKB:Q99JB6, ECO:0000269|PubMed:22678915}


FOXP3 Antibody (C-term S418) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

FOXP3 Antibody (C-term S418) - Images





FOXP3 Antibody (C-term S418)(Cat. #AP18718b) western blot analysis in CEM cell line lysates (35ug/lane). This demonstrates the FOXP3 antibody detected the FOXP3 protein (arrow).

FOXP3 Antibody (C-term S418) - Background

The protein encoded by this gene is a member of the forkhead/winged-helix family of transcriptional regulators. Defects in this gene are the cause of immunodeficiency polyendocrinopathy, enteropathy, X-linked syndrome (IPEX), also known as X-linked autoimmunity-immunodeficiency syndrome. Alternatively spliced transcript variants encoding different isoforms have been identified.

FOXP3 Antibody (C-term S418) - References

Nadeau, K., et al. J. Allergy Clin. Immunol. 126(4):845-852(2010)
Gandhi, R., et al. Nat. Immunol. 11(9):846-853(2010)
Lan, Y., et al. Zhonghua Yi Xue Yi Chuan Xue Za Zhi 27(4):433-436(2010)
Comperat, E., et al. Anal. Quant. Cytol. Histol. 32(1):39-44(2010)
Atfy, M., et al. Egypt J Immunol 16(1):117-126(2009)