

ANXA8L1/ANXA8L2 Antibody (N-term)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18762a**Specification**

ANXA8L1/ANXA8L2 Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	Q5T2P8
Other Accession	Q5VT79 , Q4FZU6 , O97529 , P13928 , NP_001092315.1
Reactivity	Human
Predicted	Rabbit, Rat
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	77-103

ANXA8L1/ANXA8L2 Antibody (N-term) - Additional Information**Other Names**

Annexin A8-like protein 1, ANXA8L1

Target/Specificity

This ANXA8L1/ANXA8L2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 77-103 amino acids from the N-terminal region of human ANXA8L1/ANXA8L2.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions


ANXA8L1/ANXA8L2 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

ANXA8L1/ANXA8L2 Antibody (N-term) - Protein Information**ANXA8L1/ANXA8L2 Antibody (N-term) - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ANXA8L1/ANXA8L2 Antibody (N-term) - Images



ANXA8L1/ANXA8L2 Antibody (N-term)(Cat. #AP18762a) western blot analysis in MDA-MB231 cell line lysates (35ug/lane). This demonstrates the ANXA8L1/ANXA8L2 antibody detected the ANXA8L1/ANXA8L2 protein (arrow).

ANXA8L1/ANXA8L2 Antibody (N-term) - Background

Annexin A8 like 2 is a member of the annexin family of evolutionarily conserved Ca^{2+} and phospholipid binding proteins. The protein may function as an anticoagulant that indirectly inhibits the thromboplastin-specific complex, and overexpression of this gene has been associated with acute myelocytic leukemia. A highly similar duplicated copy of this gene is found in close proximity on the long arm of chromosome 10.