

**RAB22A Antibody (C-term)**  
**Affinity Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP18867b****Specification**

---

**RAB22A Antibody (C-term) - Product Information**

|                   |  |
|-------------------|--|
| Application       | WB,E   |
| Primary Accession | <a href="#">O9UL26</a>                               |
| Other Accession   | <a href="#">P35285</a> , <a href="#">NP_065724.1</a> |
| Reactivity        | Human  |
| Predicted         | Mouse  |
| Host              | Rabbit   |
| Clonality         | Polyclonal   |
| Isotype           | Rabbit IgG   |
| Calculated MW     | 21855  |
| Antigen Region    | 166-192  |

**RAB22A Antibody (C-term) - Additional Information****Gene ID** 57403**Other Names**

Ras-related protein Rab-22A, Rab-22, RAB22A, RAB22

**Target/Specificity**

This RAB22A antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 166-192 amino acids from the C-terminal region of human RAB22A.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

RAB22A Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

**RAB22A Antibody (C-term) - Protein Information****Name** RAB22A

## Synonyms RAB22

**Function** Plays a role in endocytosis and intracellular protein transport. Mediates trafficking of TF from early endosomes to recycling endosomes (PubMed:[16537905](#)). Required for NGF-mediated endocytosis of NTRK1, and subsequent neurite outgrowth (PubMed:[21849477](#)). Binds GTP and GDP and has low GTPase activity. Alternates between a GTP-bound active form and a GDP-bound inactive form (PubMed:[16537905](#)).

## Cellular Location


Endosome membrane {ECO:0000250|UniProtKB:P51154}; Lipid-anchor. Cell membrane {ECO:0000250|UniProtKB:P51154}; Lipid-anchor. Early endosome. Late endosome {ECO:0000250|UniProtKB:P51154}. Cell projection, ruffle. Cytoplasmic vesicle. Cytoplasmic vesicle, phagosome. Cytoplasmic vesicle, phagosome membrane; Lipid-anchor; Cytoplasmic side. Note=Recruited to phagosomes containing S.aureus or M.tuberculosis.

## RAB22A Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

## RAB22A Antibody (C-term) - Images



RAB22A Antibody (C-term)(Cat. #AP18867b) western blot analysis in human placenta tissue lysates (35ug/lane). This demonstrates the RAB22A antibody detected the RAB22A protein (arrow).

## RAB22A Antibody (C-term) - Background

The protein encoded by this gene is a member of the RAB family of small GTPases. The GTP-bound form of the encoded protein has been shown to interact with early-endosomal antigen 1, and may be involved in the trafficking of and interaction between endosomal compartments.

**RAB22A Antibody (C-term) - References**

Zhu, H., et al. Mol. Biol. Cell 20(22):4720-4729(2009)  
Magadan, J.G., et al. Mol. Cell. Biol. 26(7):2595-2614(2006)  
Mesa, R., et al. Exp. Cell Res. 304(2):339-353(2005)  
Barrios-Rodiles, M., et al. Science 307(5715):1621-1625(2005)  
Weigert, R., et al. Mol. Biol. Cell 15(8):3758-3770(2004)