10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

FLOT1 Antibody (N-term)

Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP18905a

Specification

FLOT1 Antibody (N-term) - Product Information

Application	**WB,E**
Primary Accession	O75955
Other Accession	Q9Z1E1, Q767L6, O08917, Q08DN8, NP_005794.1
Reactivity	**Human**
Predicted	**Bovine, Mouse, Pig, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**
Isotype	**Rabbit IgG**
Calculated MW	**47355**
Antigen Region	**104-133**

FLOT1 Antibody (N-term) - Additional Information

Gene ID 10211

Other Names
Flotillin-1, FLOT1

Target/Specificity
This FLOT1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 104-133 amino acids from the N-terminal region of human FLOT1.

Dilution
WB~~1:1000

Format
Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage
Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions
FLOT1 Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

FLOT1 Antibody (N-term) - Protein Information

Name FLOT1

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999

Function May act as a scaffolding protein within caveolar membranes, functionally participating in formation of caveolae or caveolae-like vesicles.

Cellular Location
Cell membrane; Peripheral membrane protein. Endosome Membrane, caveola {ECO:0000250|UniProtKB:O08917}; Peripheral membrane protein {ECO:0000250|UniProtKB:O08917}. Melanosome. Membrane raft. Note=Identified by mass spectrometry in melanosome fractions from stage I to stage IV (PubMed:17081065) Membrane-associated protein of caveola (By similarity) {ECO:0000250|UniProtKB:O08917, ECO:0000269|PubMed:17081065}

FLOT1 Antibody (N-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

FLOT1 Antibody (N-term) - Images



FLOT1 Antibody (N-term)(Cat. #AP18905a) western blot analysis in HepG2 cell line lysates (35ug/lane).This demonstrates the FLOT1 antibody detected the FLOT1 protein (arrow).

10320 Camino Santa Fe, Suite G
San Diego, CA 92121
Tel: 858.875.1900 Fax: 858.875.1999



All lanes : Anti-FLOT1 Antibody (N-term) at 1:1000 dilution Lane 1: SK-OV-3 whole cell lysate Lane 2: Hela whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 47 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

FLOT1 Antibody (N-term) - Background

Caveolae are small domains on the inner cell membrane involved in vesicular trafficking and signal transduction. FLOT1 encodes a caveolae-associated, integral membrane protein. The function of flotillin 1 has not been determined. [provided by RefSeq].

FLOT1 Antibody (N-term) - References

Lee, B.Y., et al. Mol. Cell Proteomics 9(1):32-53(2010)
Barcellos, L.F., et al. PLoS Genet. 5 (10), E1000696 (2009) :
Saalik, P., et al. Bioconjug. Chem. 20(5):877-887(2009)
Rajendran, L., et al. PLoS ONE 4 (12), E8290 (2009) :
Rossy, J., et al. PLoS ONE 4 (4), E5403 (2009) :