

**GAS41 (YEATS4/NuBI-1) Antibody (N-term F2)**  
**Purified Rabbit Polyclonal Antibody (Pab)**  
**Catalog # AP1902d****Specification**

---

**GAS41 (YEATS4/NuBI-1) Antibody (N-term F2) - Product Information**

Application	WB,E
Primary Accession	<a href="#">O95619</a>
Other Accession	<a href="#">O9CR11</a>
Reactivity	Human
Predicted	Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	26499

**GAS41 (YEATS4/NuBI-1) Antibody (N-term F2) - Additional Information****Gene ID** 8089**Other Names**

YEATS domain-containing protein 4, Glioma-amplified sequence 41, Gas41, NuMA-binding protein 1, NuBI-1, NuBI1, YEATS4, GAS41

**Target/Specificity**

This GAS41 (YEATS4/NuBI-1) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide selected from the N-terminal region of human NuBI-1.

**Dilution**

WB~~1:1000

E~~Use at an assay dependent concentration.

**Format**

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

**Storage**

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

**Precautions**

GAS41 (YEATS4/NuBI-1) Antibody (N-term F2) is for research use only and not for use in diagnostic or therapeutic procedures.

**GAS41 (YEATS4/NuBI-1) Antibody (N-term F2) - Protein Information****Name** YEATS4 ([HGNC:24859](#))

**Function** Chromatin reader component of the NuA4 histone acetyltransferase (HAT) complex, a complex involved in transcriptional activation of select genes principally by acetylation of nucleosomal histones H4 and H2A (PubMed:[12963728](#), PubMed:[14966270](#)). Specifically recognizes and binds acylated histone H3, with a preference for histone H3 diacetylated at 'Lys-18' and 'Lys-27' (H3K18ac and H3K27ac) or histone H3 diacetylated at 'Lys-14' and 'Lys-27' (H3K14ac and H3K27ac) (PubMed:[29437725](#), PubMed:[29900004](#), PubMed:[30071723](#)). Also able to recognize and bind crotonylated histone H3 (PubMed:[30071723](#)). May also recognize and bind histone H3 succinylated at 'Lys-122' (H3K122succ); additional evidences are however required to confirm this result in vivo (PubMed:[29463709](#)). Plays a key role in histone variant H2AZ1/H2A.Z deposition into specific chromatin regions: recognizes and binds H3K14ac and H3K27ac on the promoters of actively transcribed genes and recruits NuA4-related complex to deposit H2AZ1/H2A.Z (PubMed:[29437725](#)). H2AZ1/H2A.Z deposition is required for maintenance of embryonic stem cell (By similarity).

#### Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00376, ECO:0000269|PubMed:10913114, ECO:0000269|PubMed:18445686}

#### Tissue Location


Expressed in brain, heart, kidney, liver, lung, pancreas, placenta and skeletal muscle.

### GAS41 (YEATS4/NuBI-1) Antibody (N-term F2) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### GAS41 (YEATS4/NuBI-1) Antibody (N-term F2) - Images



Anti-NuBI-1 Antibody at 1:1000 dilution + Jurkat whole cell lysate Lysates/proteins at 20 µg per

lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 26 kDa Blocking/Dilution buffer: 5% NFDM/TBST.

#### **GAS41 (YEATS4/NuBI-1) Antibody (N-term F2) - Background**

NuBI-1 is found in the nucleoli. It has high sequence homology to human MLLT1, and yeast and human MLLT3 proteins. Both MLLT1 and MLLT3 proteins belong to a class of transcription factors, indicating that the encoded protein might also represent a transcription factor. This protein is thought to be required for RNA transcription. The gene for this protein has been shown to be amplified in tumors.

#### **GAS41 (YEATS4/NuBI-1) Antibody (N-term F2) - References**

Zimmermann, K., et al., J. Biol. Chem. 277(21):18626-18631 (2002).  
Debernardi, S., et al., Blood 99(1):275-281 (2002).  
Harborth, J., et al., J. Biol. Chem. 275(41):31979-31985 (2000).  
Fischer, U., et al., Hum. Mol. Genet. 6(11):1817-1822 (1997).  
Gracia, E., et al., Hum. Mol. Genet. 5(5):595-600 (1996).