

STAG3L1 Antibody (Center)
Affinity Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP19076C**Specification**

STAG3L1 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	POCL83
Other Accession	NP_061864.2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	108-136

STAG3L1 Antibody (Center) - Additional Information**Other Names**

Putative STAG3-like protein 1, Stromal antigen 3-like protein 1, STAG3L1

Target/Specificity

This STAG3L1 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 108-136 amino acids from the Central region of human STAG3L1.

Dilution

WB~~1:2000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

STAG3L1 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

STAG3L1 Antibody (Center) - Protein Information

Name STAG3L1

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00750}.

STAG3L1 Antibody (Center) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

STAG3L1 Antibody (Center) - Images



All lanes : Anti-STAG3L1 Antibody (Center) at 1:2000 dilution Lane 1: human testis lysate Lane 2: MCF-7 whole cell lysate Lane 3: MOLT-4 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa Blocking/Dilution buffer: 5% NFDM/TBST.



All lanes : Anti-STAG3L1 Antibody (Center) at 1:2000 dilution Lane 1: MCF-7 whole cell lysate

Lane 2: ZR-75-1 whole cell lysate Lysates/proteins at 20 µg per lane. Secondary Goat Anti-Rabbit IgG, (H+L), Peroxidase conjugated at 1/10000 dilution. Predicted band size : 24 kDa
Blocking/Dilution buffer: 5% NFDM/TBST.

STAG3L1 Antibody (Center) - Background

The function of this protein remains unknown.

STAG3L1 Antibody (Center) - References

Strausberg, R.L., et al. Proc. Natl. Acad. Sci. U.S.A. 99(26):16899-16903(2002)
?.Genome Res. 8(11):1097-1108(1998)